

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C12N 15/49, 15/62, C07K 14/16, A61K 39/21</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/16884 (43) International Publication Date: 8 April 1999 (08.04.99)</p>
<p>(21) International Application Number: PCT/EP98/06040 (22) International Filing Date: 17 September 1998 (17.09.98) (30) Priority Data: 9720585.0 26 September 1997 (26.09.97) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): BRUCK, Claudine [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). GODART, Stephane, Andre, Georges [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). MARC-HAND, Martine [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). (74) Agent: TYRRELL, Arthur, William, Russell; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: FUSION PROTEINS COMPRISING HIV-1 TAT AND/OR NEF PROTEINS</p> <p>(57) Abstract</p> <p>The invention provides (a) an HIV Tat protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Nef protein or derivative thereof; or (b) an HIV Nef protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Tat protein or derivative thereof; or (c) an HIV Nef protein or derivative thereof linked to an HIV Tat protein or derivative thereof and a fusion partner. The invention further provides for a nucleic acid encoding such a protein and a host cell, such as <i>Pichia Pastoris</i>, transformed with the aforementioned nucleic acid.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

FUSION PROTEINS COMPRISING HIV-1 TAT AND/OR NEF PROTEINS

The present invention relates to novel HIV protein constructs, to their use in medicine,
5 to pharmaceutical compositions containing them and to methods of their manufacture.

In particular, the invention relates to fusion proteins comprising HIV-1 Tat and/or Nef proteins.

10 HIV-1 is the primary cause of the acquired immune deficiency syndrome (AIDS) which is regarded as one of the world's major health problems. Although extensive research throughout the world, has been conducted to produce a vaccine, such efforts thus far, have not been successful.

15 Non-envelope proteins of HIV-1 have been described and include for example internal structural proteins such as the products of the *gag* and *pol* genes and, other non-structural proteins such as Rev, Nef, Vif and Tat (Greene et al., New England J. Med, 324, 5, 308 et seq (1991) and Bryant et al. (Ed. Pizzo), *Pediatr. Infect. Dis. J.*, 11, 5, 390 et seq (1992).

20 HIV Nef and Tat proteins are early proteins, that is, they are expressed early in infection and in the absence of structural proteins.

According to the present invention there is provided a protein comprising

- 25 (a) an HIV Nef protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Tat protein or derivative thereof; or
(b) an HIV Tat protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Nef protein or derivative thereof; or
(c) an HIV Nef protein or derivative thereof linked to an HIV Tat protein or
30 derivative thereof and a fusion partner.

By 'fusion partner' is meant any protein sequence that is not Tat or Nef.

Preferably the fusion partner is protein D or its' lipidated derivative Lipoprotein D, from *Haemophilus influenzae* B. In particular, it is preferred that the N-terminal

third, i.e. approximately the first 100-130 amino acids are utilised. This is represented herein as Lipo D 1/3. In a preferred embodiment of the invention the Nef protein or derivative thereof may be linked to the Tat protein or derivative thereof. Such Nef-Tat fusions may optionally also be linked to a fusion partner, such as protein D.

5

The fusion partner is normally linked to the N-terminus of the Nef or Tat protein.

Derivatives encompassed within the present invention include molecules with a C terminal Histidine tail which preferably comprises between 5-10 Histidine residues.

10 Generally, a histidine tail containing n residues is represented herein as His (n). The presence of an histidine (or 'His') tail aids purification. More specifically, the invention provides proteins with the following structure

15	Lipo D 1/3	-	Nef	-	His (6)
	Lipo D 1/3	-	Nef-Tat	-	His (6)
	Prot D 1/3	-	Nef	-	His (6)
20	Prot D 1/3	-	Nef-Tat	-	His (6)
			Nef-Tat	-	His (6)

Figure 1 provides the amino-acid (Seq. ID. No. 7) and DNA sequence (Seq. ID. No. 6) of the fusion partner for such constructs.

25

In a preferred embodiment the proteins are expressed with a Histidine tail comprising between 5 to 10 and preferably six Histidine residues. These are advantageous in aiding purification. Separate expression, in yeast (*Saccharomyces cerevisiae*), of Nef (Macreadie I.G. et al., 1993, Yeast 9 (6) 565-573) and Tat (Braddock M et al., 1989, Cell 58 (2) 269-79) has already been reported. Nef protein only is myristilated. The present invention provides for the first time the expression of Nef and Tat separately

30

in a *Pichia* expression system (Nef-His and Tat-His constructs), and the successful expression of a fusion construct Nef-Tat-His. The DNA and amino acid sequences of representative Nef-His (Seq. ID. No.s 8 and 9), Tat-His (Seq. ID. No.s 10 and 11) and of Nef-Tat-His fusion proteins (Seq. ID. No.s 12 and 13) are set forth in Figure 2.

5

Derivatives encompassed within the present invention also include mutated proteins. The term 'mutated' is used herein to mean a molecule which has undergone deletion, addition or substitution of one or more amino acids using well known techniques for site directed mutagenesis or any other conventional method.

10

A mutated Tat is illustrated in Figure 2 (Seq. ID. No.s 22 and 23) as is a Nef-Tat Mutant-His (Seq. ID. No.s 24 and 25).

15

The present invention also provides a DNA encoding the proteins of the present invention. Such sequences can be inserted into a suitable expression vector and expressed in a suitable host.

20

A DNA sequence encoding the proteins of the present invention can be synthesized using standard DNA synthesis techniques, such as by enzymatic ligation as described by D.M. Roberts *et al.* in *Biochemistry* 1985, 24, 5090-5098, by chemical synthesis, by *in vitro* enzymatic polymerization, or by PCR technology utilising for example a heat stable polymerase, or by a combination of these techniques.

25

Enzymatic polymerisation of DNA may be carried out *in vitro* using a DNA polymerase such as DNA polymerase I (Klenow fragment) in an appropriate buffer containing the nucleoside triphosphates dATP, dCTP, dGTP and dTTP as required at a temperature of 10°-37°C, generally in a volume of 50µl or less. Enzymatic ligation of DNA fragments may be carried out using a DNA ligase such as T4 DNA ligase in an appropriate buffer, such as 0.05M Tris (pH 7.4), 0.01M MgCl₂, 0.01M dithiothreitol, 1mM spermidine, 1mM ATP and 0.1mg/ml bovine serum albumin, at a temperature of 4°C to ambient, generally in a volume of 50ml or less. The chemical synthesis of the DNA polymer or fragments may be carried out by conventional

30

phosphotriester, phosphite or phosphoramidite chemistry, using solid phase techniques such as those described in 'Chemical and Enzymatic Synthesis of Gene Fragments - A Laboratory Manual' (ed. H.G. Gassen and A. Lang), Verlag Chemie, Weinheim (1982), or in other scientific publications, for example M.J. Gait, H.W.D. Matthes, M. Singh, B.S. Sproat, and R.C. Titmas, *Nucleic Acids Research*, 1982, 10, 6243; B.S. Sproat, and W. Bannwarth, *Tetrahedron Letters*, 1983, 24, 5771; M.D. Matteucci and M.H. Caruthers, *Tetrahedron Letters*, 1980, 21, 719; M.D. Matteucci and M.H. Caruthers, *Journal of the American Chemical Society*, 1981, 103, 3185; S.P. Adams *et al.*, *Journal of the American Chemical Society*, 1983, 105, 661; N.D. Sinha, J. Biernat, J. McMannus, and H. Koester, *Nucleic Acids Research*, 1984, 12, 4539; and H.W.D. Matthes *et al.*, *EMBO Journal*, 1984, 3, 801.

The invention also provides a process for preparing a protein of the invention, the process comprising the steps of :

- i) preparing a replicable or integrating expression vector capable, in a host cell, of expressing a DNA polymer comprising a nucleotide sequence that encodes the protein or a derivative thereof
- ii) transforming a host cell with said vector
- iii) culturing said transformed host cell under conditions permitting expression of said DNA polymer to produce said protein; and
- iv) recovering said protein

The process of the invention may be performed by conventional recombinant techniques such as described in Maniatis *et al.*, *Molecular Cloning - A Laboratory Manual*; Cold Spring Harbor, 1982-1989.

The term 'transforming' is used herein to mean the introduction of foreign DNA into a host cell. This can be achieved for example by transformation, transfection or

infection with an appropriate plasmid or viral vector using e.g. conventional techniques as described in Genetic Engineering; Eds. S.M. Kingsman and A.J. Kingsman; Blackwell Scientific Publications; Oxford, England, 1988. The term 'transformed' or 'transformant' will hereafter apply to the resulting host cell
5 containing and expressing the foreign gene of interest.

The expression vectors are novel and also form part of the invention.

The replicable expression vectors may be prepared in accordance with the invention,
10 by cleaving a vector compatible with the host cell to provide a linear DNA segment having an intact replicon, and combining said linear segment with one or more DNA molecules which, together with said linear segment encode the desired product, such as the DNA polymer encoding the protein of the invention, or derivative thereof, under ligating conditions.

15

Thus, the DNA polymer may be preformed or formed during the construction of the vector, as desired.

The choice of vector will be determined in part by the host cell, which may be
20 prokaryotic or eukaryotic but preferably is *E. coli* or yeast. Suitable vectors include plasmids, bacteriophages, cosmids and recombinant viruses.

The preparation of the replicable expression vector may be carried out conventionally with appropriate enzymes for restriction, polymerisation and ligation of the DNA, by
25 procedures described in; for example, Maniatis *et al.* cited above.

The recombinant host cell is prepared, in accordance with the invention, by transforming a host cell with a replicable expression vector of the invention under transforming conditions. Suitable transforming conditions are conventional and are
30 described in, for example, Maniatis *et al.* cited above, or "DNA Cloning" Vol. II, D.M. Glover ed., IRL Press Ltd, 1985.

- The choice of transforming conditions is determined by the host cell. Thus, a bacterial host such as *E. coli* may be treated with a solution of CaCl_2 (Cohen *et al.*, Proc. Nat. Acad. Sci., 1973, 69, 2110) or with a solution comprising a mixture of RbCl , MnCl_2 , potassium acetate and glycerol, and then with 3-[N-morpholino]-propane-sulphonic acid, RbCl and glycerol. Mammalian cells in culture may be transformed by calcium co-precipitation of the vector DNA onto the cells. The invention also extends to a host cell transformed with a replicable expression vector of the invention.
- 5
- 10 Culturing the transformed host cell under conditions permitting expression of the DNA polymer is carried out conventionally, as described in, for example, Maniatis *et al.* and "DNA Cloning" cited above. Thus, preferably the cell is supplied with nutrient and cultured at a temperature below 50°C.
- 15 The product is recovered by conventional methods according to the host cell. Thus, where the host cell is bacterial, such as *E. coli* - or yeast such as *Pichia*; it may be lysed physically, chemically or enzymatically and the protein product isolated from the resulting lysate. Where the host cell is mammalian, the product may generally be isolated from the nutrient medium or from cell free extracts. Conventional protein
- 20 isolation techniques include selective precipitation, adsorption chromatography, and affinity chromatography including a monoclonal antibody affinity column.

- For proteins of the present invention provided with Histidine tails, purification can easily be achieved by the use of a metal ion affinity column. In a preferred
- 25 embodiment, the protein is further purified by subjecting it to cation ion exchange chromatography and/or Gel filtration chromatography. The protein is then sterilised by passing through a 0.22 μm membrane.

- The proteins of the invention can then be formulated as a vaccine, or the Histidine
- 30 residues enzymatically cleared.

The proteins of the present invention are provided preferably at least 80% pure more preferably 90% pure as visualised by SDS PAGE. Preferably the proteins appear as a single band by SDS PAGE.

- 5 The present invention also provides pharmaceutical composition comprising a protein of the present invention in a pharmaceutically acceptable excipient.

Vaccine preparation is generally described in **New Trends and Developments in Vaccines**, Voller *et al.* (eds.), University Park Press, Baltimore, Maryland, 1978.

- 10 Encapsulation within liposomes is described by Fullerton, US Patent 4,235,877.

- The proteins of the present invention are preferably adjuvanted in the vaccine formulation of the invention. Suitable adjuvants include an aluminium salt such as aluminium hydroxide gel (alum) or aluminium phosphate, but may also be a salt of
15 calcium, iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars, cationically or anionically derivatised polysaccharides, or polyphosphazenes.

- In the formulation of the inventions it is preferred that the adjuvant composition
20 induces a preferential TH1 response. Suitable adjuvant systems include, for example, a combination of monophosphoryl lipid A or derivative thereof, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL) together with an aluminium salt.

- An enhanced system involves the combination of a monophosphoryl lipid A and a
25 saponin derivative particularly the combination of QS21 and 3D-MPL as disclosed in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol as disclosed in WO 96/33739.

- A particularly potent adjuvant formulation involving QS21, 3D-MPL & tocopherol in
30 an oil in water emulsion is described in WO 95/17210 and is a preferred formulation.

Accordingly in one embodiment of the present invention there is provided a vaccine comprising a protein according to the invention adjuvanted with a monophosphoryl lipid A or derivative thereof, especially 3D-MPL.

- 5 Preferably the vaccine additionally comprises a saponin, more preferably QS21.

- Preferably the formulation additionally comprises an oil in water emulsion and tocopherol. The present invention also provides a method for producing a vaccine formulation comprising mixing a protein of the present invention together with a
10 pharmaceutically acceptable excipient, such as 3D-MPL.

The vaccine of the present invention may additional comprise further HIV proteins, such as the envelope glycoprotein gp160 or its derivative gp 120.

- 15 In another aspect, the invention relates to an HIV Nef or an HIV Tat protein or derivative thereof expressed in *Pichia pastoris*.

The invention will be further described by reference to the following examples:

20 **EXAMPLES:**

General

- Nef and Tat proteins, two regulatory proteins encoded by the human
25 immunodeficiency virus (HIV-1) were produced in *E.coli* and in the methylotrophic yeast *Pichia pastoris*.

- The *nef* gene from the Bru/Lai isolate (Cell 40: 9-17, 1985) was selected for these constructs since this gene is among those that are most closely related to the
30 consensus Nef.

The starting material for the Bru/Lai *nef* gene was a 1170bp DNA fragment cloned on the mammalian expression vector pcDNA3 (pcDNA3/*nef*).

The *tat* gene originates from the BH10 molecular clone. This gene was received as an HTLV III cDNA clone named pCV1 and described in Science, 229, p69-73, 1985.

1. EXPRESSION OF HIV-1 *nef* AND *tat* SEQUENCES IN E.COLI.

Sequences encoding the Nef protein as well as a fusion of *nef* and *tat* sequences were placed in plasmids vectors: pRIT14586 and pRIT14589 (see figure 1).

Nef and the Nef-Tat fusion were produced as fusion proteins using as fusion partner a part of the protein D. Protein D is an immunoglobulin D binding protein exposed at the surface of the gram-negative bacterium *Haemophilus influenzae*.

pRIT14586 contains, under the control of a λ PL promoter, a DNA sequence derived from the bacterium *Haemophilus influenzae* which codes for the first 127 amino acids of the protein D (Infect. Immun. 60 : 1336-1342, 1992), immediately followed by a multiple cloning site region plus a DNA sequence coding for one glycine, 6 histidines residues and a stop codon (Fig. 1A).

This vector is designed to express a processed lipidated His tailed fusion protein (LipoD fusion protein). The fusion protein is synthesised as a precursor with an 18 amino acid residues long signal sequence and after processing, the cysteine at position 19 in the precursor molecule becomes the amino terminal residue which is then modified by covalently bound fatty acids (Fig.1B).

pRIT14589 is almost identical to pRIT14586 except that the protD derived sequence starts immediately after the cysteine19 codon.

Expression from this vector results in a His tailed, non lipidated fusion protein (Prot D fusion protein).

Four constructs were made: LipoD-*nef*-His, LipoD-*nef-tat*-His, ProtD-*nef*-His, and ProtD-*nef-tat*-His.

The first two constructs were made using the expression vector pRIT14586, the last
5 two constructs used pRIT14589.

1.1 CONSTRUCTION OF THE RECOMBINANT STRAIN ECLD-N1 PRODUCING THE LIPOD-Nef-HIS FUSION PROTEIN.

10

1.1.1 Construction of the lipoD-*nef*-His expression plasmid pRIT14595

The *nef* gene(Bru/Lai isolate) was amplified by PCR from pcDNA3/Nef plasmid with
primers 01 and 02.

15

NcoI

PRIMER 01 (Seq ID NO 1): 5'ATCGTCCATG.GGT.GGC.AAG.TGG.T 3'

20

SpeI

PRIMER 02 (Seq ID NO 2): 5' CGGCTACTAGTGCAGTTCTTGAA 3'

The *nef* DNA region amplified starts at nucleotide 8357 and terminates at nucleotide
8971 (Cell, 40: 9-17, 1985).

25

An NcoI restriction site (which carries the ATG codon of the *nef* gene) was
introduced at the 5'end of the PCR fragment while a SpeI site was introduced at the 3'
end.

30 The PCR fragment obtained and the expression plasmid pRIT14586 were both
restricted by NcoI and SpeI, purified on an agarose gel, ligated and transformed in the

appropriate *E. coli* host cell, strain AR58. This strain is a cryptic λ lysogen derived from N99 that is *galE::Tn10*, Δ -8 (*chlD-pgl*), Δ -H1 (*cro-chlA*), N^+ , and *cI857*.

5 The resulting recombinant plasmid received, after verification of the *nef* amplified region by automatic sequencing, (see section 1.1.2 below) the pRIT14595 denomination.

1.1.2 Selection of transformants of *E. Coli* strain AR58 with pRIT14595.

10

When transformed in AR58 *E. coli* host strain, the recombinant plasmid directs the heat-inducible production of the heterologous protein.

15

Heat inducible protein production of several recombinant lipoD-Nef-His transformants was analysed by Coomassie Blue stained SDS-PAGE. All the transformants analysed showed an heat inducible heterologous protein production. The abundance of the recombinant Lipo D-Nef-Tat-His fusion protein was estimated at 10% of total protein.

20

One of the transformants was selected and given the laboratory accession number ECLD-N1.

25

The recombinant plasmid was reisolated from strain ECLD-N1, and the sequence of the *nef*-His coding region was confirmed by automated sequencing. This plasmid received the official designation pRIT14595.

The fully processed and acylated recombinant Lipo D-*nef*-His fusion protein produced by strain ECLD-N1 is composed of:

30

°Fatty acids

°109 a.a. of proteinD (starting at a.a.19 and extending to a.a.127).

°A methionine, created by the use of NcoI cloning site of pRIT14586 (Fig.1).

°205a.a. of Nef protein (starting at a.a.2 and extending to a.a.206).

5 °A threonine and a serine created by the cloning procedure (cloning at SpeI site of pRIT14586).

°One glycine and six histidines.

1.2 CONSTRUCTION OF RECOMBINANT STRAIN ECD-N1 PRODUCING PROT D-NEF-HIS FUSION PROTEIN.

10

Construction of expression plasmid pRIT14600 encoding the Prot D-Nef-His fusion protein was identical to the plasmid construction described in example 1.1.1 with the exception that pRIT14589 was used as receptor plasmid for the PCR amplified *nef* fragment.

15

E.coli AR58 strain was transformed with pRIT14600 and transformants were analysed as described in example 1.1.2. The transformant selected received laboratory accession number ECD-N1.

1.3 CONSTRUCTION OF RECOMBINANT STRAIN ECLD-NT6 PRODUCING THE LIPO D-Nef-Tat-HIS FUSION PROTEIN.

1.3.1 Construction of the lipo D-Nef-Tat-His expression plasmid pRIT14596

5

The *tat* gene(BH10 isolate) was amplified by PCR from a derivative of the pCV1 plasmid with primers 03 and 04. *SpeI* restriction sites were introduced at both ends of the PCR fragment.

10

SpeI

PRIMER 03 (Seq ID NO 3): 5' ATCGTACTAGT.GAG.CCA.GTA.GAT.C 3'

SpeI

PRIMER 04 (Seq ID NO 4): 5' CGGCTACTAGTTTTCCTTCGGGCCT 3'

15

The nucleotide sequence of the amplified *tat* gene is illustrated in the pCV1 clone (Science 229 : 69-73, 1985) and covers nucleotide 5414 till nucleotide 7998.

20 The PCR fragment obtained and the plasmid pRIT14595 (expressing lipoD-Nef-His protein) were both digested by *SpeI* restriction enzyme, purified on an agarose gel, ligated and transformed in competent AR58 cells. The resulting recombinant plasmid received, after verification of the *tat* amplified sequence by automatic sequencing (see section 1.3.2 below), the pRIT14596 denomination.

25

1.3.2 Selection of transformants of strain AR58 with pRIT14596

Transformants were grown, heat induced and their proteins were analysed by Coomassie Blue stained gels. The production level of the recombinant protein was
30 estimated at 1% of total protein. One recombinant strain was selected and received the laboratory denomination ECLD-NT6.

The lipoD-*nef*-*tat* -His recombinant plasmid was reisolated from ECLD-NT6 strain, sequenced and received the official designation pRIT14596.

5 The fully processed and acylated recombinant Lipo D-Nef-Tat-His fusion protein produced by strain ECLD-N6 is composed of:

- °Fatty acids

- °109 a.a. of proteinD (starting at a.a.19 and extending to a.a.127).

- °A methionine, created by the use of NcoI cloning site of pRIT14586.

10 °205a.a. of the Nef protein (starting at a.a.2 and extending to a.a.206)

- °A threonine and a serine created by the cloning procedure

- °85a.a. of the Tat protein (starting at a.a.2 and extending to a.a.86)

- °A threonine and a serine introduced by cloning procedure

- °One glycine and six histidines.

15

1.4 CONSTRUCTION OF RECOMBINANT STRAIN ECD-NT1 PRODUCING PROT D-Nef-Tat-HIS FUSION PROTEIN.

20 Construction of expression plasmid pRIT14601 encoding the Prot D-Nef-Tat-His fusion protein was identical to the plasmid construction described in example 1.3.1 with the exception that pRIT14600 was used as receptor plasmid for the PCR amplified *nef* fragment.

25 *E. coli* AR58 strain was transformed with pRIT14601 and transformants were analysed as described previously. The transformant selected received laboratory accession number ECD-NT1.

30

2. EXPRESSION OF HIV-1 *nef* AND *tat* SEQUENCES IN *PICHIA PASTORIS*.

Nef protein, Tat protein and the fusion Nef -Tat were expressed in the methylotrophic yeast *Pichia pastoris* under the control of the inducible alcohol oxidase (AOX1) promoter.

To express these HIV-1 genes a modified version of the integrative vector PHIL-D2 (INVITROGEN) was used. This vector was modified in such a way that expression of heterologous protein starts immediately after the native ATG codon of the AOX1 gene and will produce recombinant protein with a tail of one glycine and six histidines residues. This PHIL-D2-MOD vector was constructed by cloning an oligonucleotide linker between the adjacent *Asu*II and *Eco*RI sites of PHIL-D2 vector (see Figure 3). In addition to the His tail, this linker carries *Nco*I, *Spe*I and *Xba*I restriction sites between which *nef*, *tat* and *nef-tat* fusion were inserted.

2.1 CONSTRUCTION OF THE INTEGRATIVE VECTORS pRIT14597 (encoding Nef-His protein), pRIT14598 (encoding Tat-His protein) and pRIT14599 (encoding fusion Nef-Tat-His).

The *nef* gene was amplified by PCR from the pcDNA3/Nef plasmid with primers 01 and 02 (see section 1.1.1 construction of pRIT14595). The PCR fragment obtained and the integrative PHIL-D2-MOD vector were both restricted by *Nco*I and *Spe*I, purified on agarose gel and ligated to create the integrative plasmid pRIT14597 (see Figure 3).

The *tat* gene was amplified by PCR from a derivative of the pCV1 plasmid with primers 05 and 04 (see section 1.3.1 construction of pRIT14596):

*Nco*I

PRIMER 05 (Seq ID NO 5): 5'ATCGTCCATGGAGCCAGTAGATC 3'

An NcoI restriction site was introduced at the 5' end of the PCR fragment while a SpeI site was introduced at the 3' end with primer 04. The PCR fragment obtained and the PHIL-D2-MOD vector were both restricted by NcoI and SpeI, purified on agarose gel and ligated to create the integrative plasmid pRIT14598.

5

To construct pRIT14599, a 910bp DNA fragment corresponding to the *nef-tat*-His coding sequence was ligated between the EcoRI blunted(T4 polymerase) and NcoI sites of the PHIL-D2-MOD vector. The *nef-tat*-His coding fragment was obtained by XbaI blunted(T4 polymerase) and NcoI digestions of pRIT14596.

10

2.2 TRANSFORMATION OF PICHIA PASTORIS STRAIN GS115(his4).

To obtain *Pichia pastoris* strains expressing Nef-His, Tat-His and the fusion Nef-Tat-His, strain GS115 was transformed with linear NotI fragments carrying the respective expression cassettes plus the HIS4 gene to complement his4 in the host genome. Transformation of GS115 with NotI-linear fragments favors recombination at the AOXI locus.

15

Multicopy integrant clones were selected by quantitative dot blot analysis and the type of integration, insertion (Mut⁺ phenotype) or transplacement (Mut⁺ phenotype), was determined.

20

From each transformation, one transformant showing a high production level for the recombinant protein was selected :

25

Strain Y1738 (Mut⁺ phenotype) producing the recombinant Nef-His protein, a myristylated 215 amino acids protein which is composed of:

30

°Myristic acid

°A methionine, created by the use of NcoI cloning site of PHIL-D2-MOD vector

°205 a.a. of Nef protein(starting at a.a.2 and extending to a.a.206)

°A threonine and a serine created by the cloning procedure (cloning at SpeI site of PHIL-D2-MOD vector.

°One glycine and six histidines.

- 5 Strain Y1739 (Mut⁺ phenotype) producing the Tat-His protein, a 95 amino acid protein which is composed of:

°A methionine created by the use of NcoI cloning site

°85 a.a. of the Tat protein(starting at a.a.2 and extending to a.a.86)

10

°A threonine and a serine introduced by cloning procedure

°One glycine and six histidines

- Strain Y1737(Mut⁺ phenotype) producing the recombinant Nef-Tat-His fusion protein,
15 a myristylated 302 amino acids protein which is composed of:

°Myristic acid

°A methionine, created by the use of NcoI cloning site

°205a.a. of Nef protein(starting at a.a.2 and extending to a.a.206)

20

°A threonine and a serine created by the cloning procedure

°85a.a. of the Tat protein(starting at a.a.2 and extending to a.a.86)

°A threonine and a serine introduced by the cloning procedure

°One glycine and six histidines

3. EXPRESSION OF HIV-1 Tat-MUTANT IN PICHIA PASTÓRIS

As well as a Nef-Tat mutant fusion protein, a mutant recombinant Tat protein has also
5 been expressed. The mutant Tat protein must be **biologically inactive** while
maintaining its immunogenic epitopes.

A double mutant *tat* gene, constructed by D.Clements (Tulane University) was
selected for these constructs.

10

This *tat* gene (originates from BH10 molecular clone) bears **mutations** in the **active
site region (Lys41→Ala)** and in **RGD motif (Arg78→Lys and Asp80→Glu)** (
Virology 235: 48-64, 1997).

15

The mutant *tat* gene was received as a cDNA fragment subcloned between the EcoRI
and HindIII sites within a CMV expression plasmid (pCMVLys41/KGE)

3.1 CONSTRUCTION OF THE INTEGRATIVE VECTORS

20

**pRIT14912(encoding Tat mutant-His protein) and pRIT14913(encoding fusion
Nef-Tat mutant-His).**

The *tat* mutant gene was amplified by PCR from the pCMVLys41/KGE plasmid with
primers 05 and 04 (see section 2.1 construction of pRIT14598)

25

An NcoI restriction site was introduced at the 5' end of the PCR fragment while a
SpeI site was introduced at the 3' end with primer 04. The PCR fragment obtained and
the PHIL-D2-MOD vector were both restricted by NcoI and SpeI, purified on agarose
gel and ligated to create the integrative plasmid pRIT14912

30

To construct pRIT14913, the *tat* mutant gene was amplified by PCR from the pCMVLys41/KGE plasmid with primers 03 and 04 (see section 1.3.1 construction of pRIT14596).

- 5 The PCR fragment obtained and the plasmid pRIT14597 (expressing Nef-His protein) were both digested by SpeI restriction enzyme, purified on agarose gel and ligated to create the integrative plasmid pRIT14913

3.2 TRANSFORMATION OF PICHIA PASTORIS STRAIN GS115.

10

Pichia pastoris strains expressing Tat mutant-His protein and the fusion Nef-Tat mutant-His were obtained, by applying integration and recombinant strain selection strategies previously described in section 2.2 .

- 15 Two recombinant strains producing Tat mutant-His protein ,a 95 amino-acids protein, were selected: Y1775 (Mut⁺ phenotype) and Y1776(Mut^s phenotype).

One recombinant strain expressing Nef-Tat mutant-His fusion protein, a 302 amino-acids protein was selected: Y1774(Mut⁺ phenotype).

20

4. PURIFICATION OF Nef-Tat-His FUSION PROTEIN (PICHIA PASTORIS)

- 5 The purification scheme has been developed from 146g of recombinant Pichia pastoris cells (wet weight) or 2L Dyno-mill homogenate OD 55. The chromatographic steps are performed at room temperature. Between steps, Nef-Tat positive fractions are kept overnight in the cold room (+4°C); for longer time, samples are frozen at -20°C.

10

146g of Pichia pastoris cells



Homogenization

Buffer: 2L 50 mM PO₄ pH 7.0

final OD:50



Dyno-mill disruption (4 passes)



Centrifugation

JA10 rotor / 9500 rpm/ 30 min / room temperature



Dyno-mill Pellet



Wash

(1h - 4°C)

Buffer: +2L 10 mM PO₄ pH 7.5 -

150mM - NaCl 0,5% empigen



Centrifugation

JA10 rotor / 9500 rpm/ 30 min / room temperature



Pellet

↓
Solubilisation

(O/N - 4°C)

Buffer: + 660ml 10 mM PO₄ pH 7.5 -
150mM NaCl - 4.0M GuHCl

↓
Reduction

(4H - room temperature - in the dark)

+ 0,2M 2-mercaptoethanesulfonic acid,
sodium salt (powder addition) / pH
adjusted to 7.5 (with 0,5M NaOH
solution) before incubation

↓
Carboxymethylation

(1/2 h - room temperature - in the dark)

+ 0,25M Iodoacetamid (powder addition)
/ pH adjusted to 7.5 (with 0,5M NaOH
solution) before incubation

↓
Immobilized metal ion affinity chromatography on
Ni²⁺-NTA-Agarose (Qiagen - 30 ml of resin)

Equilibration buffer: 10 mM PO₄ pH 7.5 -
150mM NaCl - 4.0M GuHCl

Washing buffer: 1) Equilibration
buffer

2) 10 mM PO₄ pH
7.5 - 150mM NaCl - 6M Urea

3) 10 mM PO₄ pH
7.5 - 150mM NaCl - 6M Urea - 25
mM Imidazol

Elution buffer: 10 mM PO₄ pH 7.5 -
150mM NaCl - 6M Urea - 0,5M Imidazol


↓
Dilution

Down to an ionic strength of 18 mS/cm²

Dilution buffer: 10 mM PO₄ pH 7.5 - 6M
Urea

↓
Cation exchange chromatography on SP Sepharose FF
(Pharmacia - 30 ml of resin)

Equilibration buffer: 10 mM PO₄ pH 7.5
- 150mM NaCl - 6.0M Urea



* ratio: 0,5M Arginin for a protein concentration of 1600µg/ml.

5 Purity

The level of purity as estimated by SDS-PAGE is shown in Figure 4 by Daiichi Silver Staining and in Figure 5 by Coomassie blue G250.

After Superdex200 step: > 95%
After dialysis and sterile filtration steps: > 95%

5 Recovery

51mg of Nef-Tat-his protein are purified from 146g of recombinant *Pichia pastoris* cells (= 2L of Dyno-mill homogenate OD 55)

10 5. VACCINE PREPARATION

A vaccine prepared in accordance with the invention comprises the expression product of a DNA recombinant encoding an antigen as exemplified in example 1 or 2 and as adjuvant, the formulation comprising a mixture of 3 de -O-acylated monophosphoryl
15 lipid A 3D-MPL and QS21 in an oil/water emulsion.

3D-MPL: is a chemically detoxified form of the lipopolysaccharide (LPS) of the Gram-negative bacteria *Salmonella minnesota*.

20 Experiments performed at Smith Kline Beecham Biologicals have shown that 3D-MPL combined with various vehicles strongly enhances both the humoral and a TH1 type of cellular immunity.

QS21: is one saponin purified from a crude extract of the bark of the *Quillaja Saponaria* Molina tree, which has a strong adjuvant activity: it activates both antigen-specific lymphoproliferation and CTLs to several antigens.
25 Experiments performed at Smith Kline Beecham Biologicals have demonstrated a clear synergistic effect of combinations of 3D-MPL and QS21 in the induction of both humoral and TH1 type cellular immune responses.

30

The oil/water emulsion is composed of 2 oils (a tocopherol and squalene), and of PBS containing Tween 80 as emulsifier. The emulsion comprised 5% squalene 5%

tocopherol 0.4% Tween 80 and had an average particle size of 180 nm (see WO 95/17210).

- 5 Experiments performed at Smith Kline Beecham Biologicals have proven that the adjunction of this O/W emulsion to 3D-MPL/QS21 further increases their immunostimulant properties.

Preparation of the oil/water emulsion (2 fold concentrate)

- 10 Tween 80 is dissolved in phosphate buffered saline (PBS) to give a 2% solution in the PBS. To provide 100ml two fold concentrate emulsion 5g of DL alpha tocopherol and 5ml of squalene are vortexed to mix thoroughly. 90ml of PBS/Tween solution is added and mixed thoroughly. The resulting emulsion is then passed through a syringe and finally microfluidised by using an M110S microfluidics machine. The resulting
15 oil droplets have a size of approximately 180 nm.

Preparation of oil in water formulation.

- Antigen prepared in accordance with example 1 or 2 (5µg) was diluted in 10 fold
20 concentrated PBS pH 6.8 and H₂O before consecutive addition of SB62, 3D-MPL (5µg), QS21 (5µg) and 50 µg/ml thiomersal as preservative at 5 min interval. The emulsion volume is equal to 50% of the total volume (50µl for a dose of 100µl).

- All incubations were carried out at room temperature with agitation.
25

6. IMMUNOGENICITY OF Tat AND Nef-Tat IN RODENTS

- Characterization of the immune response induced after immunization with Tat and
30 NefTat was carried out. To obtain information on isotype profiles and cell-mediated immunity (CMI) two immunization experiments in mice were conducted. In the first experiment mice were immunized twice two weeks apart into the footpad with Tat or

- NefTat in the oxydized or reduced form, respectively. Antigens were formulated in an oil in water emulsion comprising squalene, tween 80[™] (polyoxyethylene sorbitan monooleate) QS21, 3D-MPL and α -tocopherol, and a control group received the adjuvant alone. Two weeks after the last immunization sera were obtained and
- 5 subjected to Tat-specific ELISA (using reduced Tat for coating) for the determination of antibody titers and isotypes (Figure 6a). The antibody titers were highest in the mice having received oxydized Tat. In general, the oxydized molecules induced higher antibody titers than the reduced forms, and Tat alone induced higher antibody titers than NefTat. The latter observation was confirmed in the second experiment.
- 10 Most interestingly, the isotype profile of Tat-specific antibodies differed depending on the antigens used for immunization. Tat alone elicited a balanced IgG1 and IgG2a profile, while NefTat induced a much stronger T_{H2} bias (Figure 6b). This was again confirmed in the second experiment.
- 15 In the second mouse experiment animals received only the reduced forms of the molecules or the adjuvant alone. Besides serological analysis (see above) lymphoproliferative responses from lymph node cells were evaluated. After restimulation of those cells in vitro with Tat or NefTat ³H-thymidine incorporation was measured after 4 days of culture. Presentation of the results as stimulation indices
- 20 indicates that very strong responses were induced in both groups of mice having received antigen (Figure 7).

In conclusion, the mice studies indicate that Tat as well as Nef-Tat are highly immunogenic candidate vaccine antigens. The immune response directed against the

25 two molecules is characterized by high antibody responses with at least 50% IgG1. Furthermore, strong CMI responses (as measured by lymphoproliferation) were observed.

7. FUNCTIONAL PROPERTIES OF THE Tat AND Nef-Tat PROTEINS

30

The Tat and NefTat molecules in oxydized or reduced form were investigated for their ability to bind to human T cell lines. Furthermore, the effect on growth of

those cell lines was assessed. ELISA plates were coated overnight with different concentration of the Tat and NefTat proteins, the irrelevant gD from herpes simplex virus type II, or with a buffer control alone. After removal of the coating solution HUT-78 cells were added to the wells. After two hours of incubation the wells were
5 washed and binding of cells to the bottom of the wells was assessed microscopically. As a quantitative measure cells were stained with toluidine blue, lysed by SDS, and the toluidine blue concentration in the supernatant was determined with an ELISA plate reader. The results indicate that all four proteins, Tat and NefTat in oxydized or reduced form mediated binding of the cells to the
10 ELISA plate (Figure 8). The irrelevant protein (data not shown) and the buffer did not fix the cells. This indicates that the recombinantly expressed Tat-containing proteins bind specifically to human T cell lines.

In a second experiment HUT-78 cells were left in contact with the proteins for 16
15 hours. At the end of the incubation period the cells were labeled with [^3H]-thymidine and the incorporation rate was determined as a measure of cell growth. All four proteins included in this assay inhibited cell growth as judged by diminished radioactivity incorporation (Figure 9). The buffer control did not mediate this effect. These results demonstrate that the recombinant Tat-containing
20 proteins are capable of inhibiting growth of a human T cell line.


In summary the functional characterization of the Tat and NefTat proteins reveals that these proteins are able to bind to human Tcell lines. Furthermore, the proteins are able to inhibit growth of such cell lines.

CLAIMS

1. A protein comprising
 - 5 (a) an HIV Tat protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Nef protein or derivative thereof; or
 - (b) an HIV Nef protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Tat protein or derivative thereof; or
 - (c) an HIV Nef protein or derivative thereof linked to an HIV Tat protein or
 - 10 derivative thereof and a fusion partner;
2. A protein as claimed in claim 1 which is a Tat-Nef fusion protein or derivative thereof.
- 15 3. A protein as claimed in claim 1 which is a Nef-Tat fusion protein or derivative thereof.
4. A protein according to claim 1 wherein the derivative of the Tat protein is a
- 20 mutated Tat protein.
5. A protein according to claim 1 wherein the derivative of the Nef protein is a mutated Nef protein.
- 25 6. A Protein as claimed in any one of claims 1 - 5 wherein the fusion partner is a lipoprotein or derivative thereof.
7. A protein as claimed in claim 6 wherein the lipoprotein is Haemophilus Influenza B protein D or derivative thereof.
- 30

8. A protein as claimed in Claim 7 wherein the fusion partner comprises between 100-130 amino acid from the N terminal of Haemophilus Influenza B protein D.
- 5 9. A protein as claimed in any one of Claims 1 to 8, wherein the Tat protein is the entire Tat protein.
- 10 10. A protein as claimed in any one of Claims 1 to 8, wherein the Nef protein is the entire Nef protein.
11. A protein as claimed in any one of Claims 1 to 10, wherein the Tat protein is fused to an HIV Nef protein and a fusion partner.
- 15 12. A protein as claimed in any one of Claims 1 to 11, wherein the protein has a Histidine tail.
13. A nucleic acid encoding a protein of Claims 1 to 12.
14. A host transformed with a nucleic acid of Claim 13.
- 20 15. A host as claimed in claim 14 wherein the host is either Pichia pastoris or E. coli.
- 25 16. A vaccine comprising a protein of any one of Claims 1 to 12 in admixture with a pharmaceutically acceptable excipient.
17. A vaccine of Claim 16 additionally comprising an adjuvant.
- 30 18. A vaccine of claim 17 wherein the adjuvant is a TH1 inducing adjuvant.

19. A vaccine as claimed in Claim 17 or 18 which adjuvant comprises monophosphoryl lipid A or derivative thereof such as 3 de-O-acylated monophosphoryl lipid A.
- 5 20. A vaccine as claimed in any one of Claims 16 to 19 additionally comprising a saponin adjuvant.
21. A method of producing a protein of Claim 1 to 12, comprising the steps of transforming a host with a nucleic acid encoding said protein, expressing said protein and recovering the protein.
- 10 22. A method as claimed in Claim 21 wherein the host is *E. coli.* or *Pichia pastoris.*
- 15 23. A method of producing a vaccine of Claim 16 to 20, comprising admixing the protein of Claim 1 to 12 with a pharmaceutically acceptable diluent.
24. A method of preparing (i) an HIV Nef protein or derivative thereof or (ii) an HIV Tat protein or derivative thereof in *Pichia pastoris* which method comprises the steps of transforming *Pichia pastoris* with DNA encoding said HIV Nef protein or derivative thereof or HIV Tat protein or derivative thereof, expressing said protein and recovering the protein.
- 20
- 25
- 30

Figure 1: A/ Map of plasmid pRIT14586

B/ Coding sequence of the first 127 amino acids
of protein D and multiple cloning site. The signal
sequence is underlined.

BamHI
ATG GAT CCA AAA ACT TTA GCC CTT TCT TTA TTA GCA GCT GGC GTA CTA GCA GGT TGT AGC AGC
 Met Asp Pro Lys Thr Leu Ala Leu Ser Leu Leu Ala Ala Gly Val Leu Ala Gly Cys Ser Ser
 CAT TCA TCA AAT ATG GCG AAT ACC CAA ATG AAA TCA GAC AAA ATC ATT ATT GCT CAC CGT GGT
 His Ser Ser Asn Met Ala Asn Thr Gln Met Lys Ser Asp Lys Ile Ile Ile Ala His Arg Gly
 GCT AGC GGT TAT TTA CCA GAG CAT ACG TTA GAA TCT AAA GCA CTT GCT TTT GCA CAA CAG GCT
 Ala Ser Gly Tyr Leu Pro Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala
 GAT TAT TTA GAG CAA GAT TTA GCA ATG ACT AAG GAT GGT CGT TTA GTG GTT ATT CAC GAT CAC
 Asp Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val Ile His Asp His
 TTT TTA GAT GGC TTG ACT GAT GTT GCG AAA AAA TTC CCA CAT CGT CAT CGT AAA GAT GGC CGT
 Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe Pro His Arg His Arg Lys Asp Gly Arg
 TACTAT GTC ATC GAC TTT ACC TTA AAA GAA ATT GAA AGT TTA GAA ATG ACA GAA AAC TTT GAA
 Tyr Tyr Val Ile Asp Phe Thr Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu
NcoI ACC ATG GCC ACG TGT GAT CAG AGC TCA SpeI ACT AGT GGA CAC CAT CAC CAT CAC CAT TAA TCT XbaI AGA
 Thr Met Ala Thr Cys Asp Gln Ser Ser Thr Ser Gly His His His His His His •

The amino acid sequence of Figure 1 relates to Seq. ID no. 7 and the nucleic acid sequence of
Figure 1 relates to Seq. ID. No. 6.

The DNA and amino acid sequences of Nef-His; Tat-His; Nef-Tat-His fusion and mutated Tat is illustrated.

Pichia-expressed constructs (plain constructs)

⇒ Nef - HIS

DNA sequence (Seq. ID. No. 8)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGA
ATGAGACGAGCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAA
AAACATGGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGG
CTAGAAGCACAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTA
AGACCAATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAGAAAAGGGG
GGACTGGAAGGGCTAATTCACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATC
TACCACACACAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTC
AGATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAG
GTAGAAGAGGCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCAT
GGAATGGATGACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCA
TTTCATCACGTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGC
CACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 9)

MGGKWSKSSVVGWPTVRRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAW
LEAQEEEEVGFVPTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWI
YHTQGYFPDWQNYTPGPGVRYPLTFGWCYKLPVPEPDKVEEANKGENTSLLHPVSLH
GMDDPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSGHHHHHH.

⇒ Tat - HIS

DNA sequence (Seq. ID. No. 10)

ATGGAGCCAGTAGATCCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAA
ACTGCTTGTACCAATTGCTATTGTAAAAAGTGTGCTTTTCATTGCCAAGTTTGTTC
ATAACAAAAGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGA
CCTCCTCAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAA

TCCCGAGGGGACCCGACAGGCCCGAAGGAACTAGTGGCCACCATCACCATCACCAT
TAA

Protein sequence (Seq. ID. No. 11)

MEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFHCQVCFITKALGISYGRKKRRQRRR
PPQGSQTHQVSLSKQPTSQSRGDPTGPKETSGHHHHHH.

⇒ Nef - Tat - HIS

DNA sequence (Seq. ID. No. 12)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGA
ATGAGACGAGCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAA
AAACATGGAGCAATCACAAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGG
CTAGAAGCACAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTA
AGACCAATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGG
GGACTGGAAGGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATC
TACCACACACAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTC
AGATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAG
GTAGAAGAGGCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCAT
GGAATGGATGACCCTGAGAGAGAAGTGTAGAGTGGAGGTTTGACAGCCGCTAGCA
TTTCATCACGTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAG
CCAGTAGATCCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAAGTGTCT
TGTACCAATTGCTATTGTAAAAAGTGTGCTTTCATTGCCAAGTTTGTTCATAACA
AAAGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCT
CAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGA
GGGGACCCGACAGGCCCGAAGGAAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 13)

^^

MGGKWSKSSVVGWPTVRRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAW
LEAQEEEEVGFVPVTPQVPLRPMYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWI
YHTQGYFPDWQNYTPGPGVRYPLTFGWCYKLVPVEPDKVEEANKGENTSLLHPVSLH
GMDDPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTA
CTNCYCKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSR
GDPTGPKETSGHHHHHH.

E.coli-expressed constructs (fusion constructs)

⇒ LipoD-Nef-HIS

DNA sequence (Seq. ID. No. 14)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

The Lipidation Signal Sequence is underlined. After processing, the cysteine coded by the TGT codon, indicated with a star, becomes the amino terminal residue which is then modified by covalently bound fatty acids.

*

ATGGATCCAAAACTTTAGCCCTTTCTTTATTAGCAGCTGGCGTACTAGCAGGTTGT
 AGCAGCCATTTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAAATCATTATT
 GCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCTAAAGCACTT
 GCTTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACTAAGGATGGT
 CGTTTAGTGTTATTACGATCACTTTTTAGATGGCTTGACTGATGTTGCGAAAAAA
 TTCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTTACCTTAAAA
 GAAATTCAAAGTTTAGAAATGACAGAAAACCTTTGAAACCATGGGTGGCAAGTGGTCA
 AAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCA
 GCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGAGCAATCACA
 AGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCACAAGAGGAG
 GAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACTTACAAG
 GCAGCTGTAGATCTTAGCCACTTTTTTAAAGAAAAGGGGGGACTGGAAGGGCTAATT
 CACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC
 TTCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTCAGATATCCACTGACCTTT
 GGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAA
 GGAGAGAACACCAGCTTGTTACACCTGTGAGCCTGCATGGAATGGATGACCCTGAG
 AGAGAAGTGTTAGAGTGAGGTTTGACAGCCGCCTAGCATTTTCATCACGTGGCCCGA
 GAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGCCACCATCACCATCACCAT
 TAA

Protein sequence of the processed lipidated ProtD-Nef-HIS protein (Seq. ID. No. 15)

(Amino-acids corresponding to Prot D fusion partner are in bold)

CSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMTKD
 GRLVVIHDHFLDGLTDVAKKFPHRHRKDGRYYVIDFTLKEIQSLEMTENFETMGGKW
 SKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEAQE
 EEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWIYHTQG
 YFPDWQNYTPGPGVRYPLTFGWICYKLVPVEPDKVEEANKGENTSLHHPVSLHGMDDP
 EREVLEWRFDSRLAFHHVARELHPEYFKNCTSGHHHHHHH.

⇒ LipoD-Nef-Tat-HIS

DNA sequence (Seq. ID. No. 16)

5/17

Nucleotides corresponding to the Prot D Fusion Partner are in bold.
The Lipidation Signal Sequence is underlined. After processing, the cysteine coded by the TGT codon, indicated with a star, becomes the amino terminal residue which is then modified by covalently bound fatty acids.

*

ATGGATCCAAAACTTTAGCCCTTTCTTTATTAGCAGCTGGCGTACTAGCAGGTTGT
AGCAGCCATTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAAATCATTATT
GCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATAACGTTAGAATCTAAAGCACTT
GCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACTAAGGATGGT
CGTTTAGTGGTTATTCACGATCACTTTTTAGATGGCTTGACTGATGTTGCGAAAAAA
TTCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTTACCTTAAAA
GAAATTCAAAGTTT**AGAAATGACAGAAAAC****TTTGAAACC****ATGGGTGGCAAGTGGTCA**
AAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCA
GCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGAGCAATCACA
AGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCACAAGAGGAG
GAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACTTACAAG
GCAGCTGTAGATCTTAGCCACTTTTTAAAGAAAAGGGGGGACTGGAAGGGGCTAATT
CACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC
TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCACTGACCTTT
GGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAA
GGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGATGACCCTGAG
AGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCACGTGGCCCGA
GAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGATCCTAGACTA
GAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAACTGCTTGTACCAATTGCTATTGT
AAAAAGTGTTGCTTTTATTGCCAAGTTTGTTCATAACAAAAGCCTTAGGCATCTCC
TATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCAAGGCAGTCAGACTCAT
CAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGAGGGGACCCGACAGGCCCG
AAGGAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence of the processed lipidated ProtD-NEF-TAT-HIS protein (Seq. ID. No. 17)

(Amino-acids corresponding to Prot D fusion partner are in bold)

CSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMTKD
GRLVVIHDHFLDGLTDVAKKFPHRHRKDGRYYVIDFTLKEIQSLEMTENFETMGGKW
SKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEAQE
EEEVGFVPTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWLYHTQG
YFPDWQNYTPGPGVRYPLTFGWCYKLVPVEPDKVEEANKGENTSLLHPVSLHGMDDP
EREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLPEWKHPGSQPKTACTNCY
CKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSRGDPTG
PKETSGHHHHHHH.

6/17

⇒ ProtD-Nef -HISDNA sequence (Seq. ID. No. 18)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

ATGGATCCAAGCAGCCATTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAA
ATCATTATTGCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATAACGTTAGAATCT
AAAGCACTTGCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACT
AAGGATGGTCGTTTGTAGTGGTTATTCACGATCACTTTTGTAGATGGCTTGACTGATGTT
GCGAAAAAATCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTT
ACCTTAAAAGAAATTCAAAGTTTAGAAATGACAGAAAACTTTGAAACCATGGGTGGC
AAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGA
GCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGA
GCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCCTGGCTAGAAGCA
CAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATG
ACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTAAAAGAAAAGGGGGGACTGGAA
GGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACA
CAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCA
CTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAG
GCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGAT
GACCCTGAGAGAGAAAGTGTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCAC
GTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGCCACCATCAC
CATCACCATTAA

Protein sequence (Seq. ID. No. 19)

(Amino-acids corresponding to Prot D fusion partner are in bold)

MDPSSHSSNMANTQMKS~~DKII~~AHRGASGYLPEHTLESKALAF~~QQADYL~~
EQDLAMTKD~~GRLVVI~~HDHFLDGLTDVAKK~~FPHRHR~~KDGRYYVIDFTLK
EIQSLEM~~TENFET~~MG~~GKWS~~KSSVVGWPTVRERM~~RAEPAADGVGAAS~~RDL
EKHGAI~~TSSNTAATNAACA~~WLEAQEEEEVGFPVTPQVPLRPMTYKA~~AVDLSH~~
FLKEKGGLEGLIHSQRRQDILDLWYHTQGYFPDWQNYTPGPGVRYPLTFGW
CYKLVPVEPDKVEEANKGENTSLLHPVSLHGMDDPEREVLEWRFD~~SRLAFH~~
HVARELHPEYFKNCTSGHHHHHH.

⇒ ProtD-Nef -Tat-HISDNA sequence (Seq. ID. No. 20)

7/17

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

ATGGATCCAAGCAGCCATTTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAA
ATCATTATTGCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCT
AAAGCACTTGCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACT
AAGGATGGTCGTTTGTAGTGGTTATTCACGATCACTTTTGTAGATGGCTTGACTGATGTT
GCGAAAAAATTCCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTT
ACCTTAAAAGAAATTCAAAGTTTAGAAATGACAGAAAACTTTGAAACCATGGGTGGC
AAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGA
GCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGA
GCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCA
CAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATG
ACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGGGGACTGGAA
GGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACA
CAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTCAGATATCCA
CTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAG
GCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGAT
GACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCAC
GTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGAT
CCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAACCTGCTTGTACCAAT
TGCTATTGTAAAAAGTGTTGCTTTTCATTGCCAAGTTTGTTCATAACAAAAGCCTTA
GGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCAAGGCAGT
CAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGAGGGGACCCG
ACAGGCCCCGAAGGAAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 21)

(Amino-acids corresponding to Prot D fusion partner are in bold)

MDPSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMT
KDGRLLVVIHDFLDGLTDVAKKFPHRHRKDGRYYVIDFTLKEIQSLEMTENFETMGG
KWSKSSVVGWPTVRRMRRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEA
QEEEEVGFVPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWYIYHT
QGYFPDWQNYTPGPGVRYPLTFGWICYKLVPEPDKVEEANKGENTSLLHPVSLHGMD
DPEREVLEWRFDLSRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTACTN
CYCKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSRGDP
TGPKETSGHHHHHH.

⇒ Tat-MUTANT-HIS

DNA sequence (Seq. ID. No. 22)

ATGGAGCCAGTAGATCCTAGACTAGAGCCCTGGAAGCATC 40
CAGGAAGTCAGCCTAAACTGCTTGTACCAATTGCTATTG 80
TAAAAAGTGTGCTTTTCATTGCCAAGTTTGTTCATAACA 120
GCTGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGAC 160
AGCGACGAAGACCTCCTCAAGGCAGTCAGACTCATCAAGT 200
TTCTCTATCAAAGCAACCCACCTCCCAATCCAAAGGGGAG 240
CCGACAGGCCCGAAGGAACTAGTGGCCACCATCACCATC 280
ACCATTAA 288

Protein sequence(Seq. ID. No. 23)

Mutated amino-acids in Tat sequences are in bold.

MEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFHCQVCFIT 40
AALGISYGRKKRRRQRRRPQGSQTHQVSLSKQPTSQSKGE 80
PTGPKETSGHHHHHH. 95

⇒Nef-Tat-Mutant-HIS

DNA sequence(Seq. ID. No. 24)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGC 40
CTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCAGCAGC 80
AGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACAT 120
GGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTG 160
CTTGTGCCTGGCTAGAAGCACAAGAGGAGGAGGAGGTGGG 200
TTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACT 240
TACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAGAAA 280
AGGGGGGACTGGAAGGGCTAATCACTCCCAACGAAGACA 320
AGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC 360
TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTCA 400
GATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACC 440
AGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAAGGAGAG 480
AACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGG 520
ATGACCCTGAGAGAGAAGTGTAGAGTGGAGGTTTGACAG 560
CCGCCTAGCATTTTCATCACGTGGCCCGAGAGCTGCATCCG 600
GAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGATCCTA 640
GACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAC 680
TGCTTGTAACCAATTGCTATTGTAAAAAGTGTGCTTTCAT 720
TGCCAAGTTTGTTCATAACAGCTGCCTTAGGCATCTCCT 760
ATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCA 800
AGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCC 840
ACCTCCCAATCCAAAGGGGAGCCGACAGGCCCGAAGGAAA 880
CTAGTGGCCACCATCACCATCACCATTAA 909


9/17

Protein sequence (Seq. ID. No. 25)

Mutated amino-acids in Tat sequence are in bold.

MGGKWSKSSVVGWPTVRERMRAEPAADGVGAASRDLEKH 40
GAITSSNTAATNAACAWLEAQEEEEVGFPVTPQVPLRPMT 80
YKAAVDLSHFLKEKGGLEGLIHSQRRQDILDLWIYHTQGY 120
FPDWQNYTPGPGVRYPLTFGWCYKLVPVEPDKVEEANKGE 160
NTSL LHPVSLHGMDDPEREVLEWRFD SRLAFHHVARELHP 200
EYFKNCTSEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFH 240
CQVCFITAALGISYGRKKRRQRRRPPQGSQTHQVSLSKQP 280
TSQSKGEPTGPKETSGHHHHHH. 302

Fig. 3 Map of pRIT14597 integrative vector




MCS POLYLINKER: *nef* gene inserted between *NcoI* and *SpeI* sites.

<i>Acu II</i>	<i>Nco I</i>	<i>Spe I</i>	<i>Eco RI</i>
TTCGAA	ACC.ATGGCCGGGACTAGT	GGC.CAC.CAT.CAC.CAT.CAC.CAT.TAA	CGGAATTC
	Thr . Ser . Gly .	His . His . His . His . His . His	

The amino acid sequence of Figure 3 relates to Seq. ID no. 27 and the nucleic acid sequence of Figure 3 relates to Seq. ID. No.26.

Fig . 4 SDS-PAGE: Nef-Tat-his fusion protein

- 1: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa
- 2: TNH/23 SP eluate (250 ng)
- 3: TNH/23 Purified bulk (250 ng)
- 4: TNH/22 Purified bulk (250 ng)
- 5: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa
- 6: TNH/23 SP eluate (400 ng)
- 7: TNH/23 Purified bulk (400 ng)
- 8: TNH/22 Purified bulk (400 ng)



Daiichi Silver Staining




Blot Tat2



BlotαNef-Tat (LAS 97340)

12/17

Fig . 5 SDS-PAGE: Nef-Tat-his fusion proteinCoomassie blue G250

- 1: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa)
- 2: TNH/23 SP eluate (4 µg)
- 3: TNH/23 Superdex200 eluate (4 µg)
- 4: TNH/23 Purified bulk (4 µg)
- 5: TNH/22 Purified bulk (4 µg)

- 6: TNH/23 Purified bulk (4 µg) / non reducing conditions
- 7: TNH/22 Purified bulk (4 µg) / non reducing conditions

Fig. 6A Tat-specific antibody titers and isotypes

group	immunization	midpoint titers					ratio IgG1/IgG2a
		Ig	IgG1	IgG2a	IgG2b		
1	oxydized Tat	353557	135538	98771	98763	1,372	
2	reduced Tat	252275	72087	76273	72014	0,945	
3	oxydized Nef-Tat	246466	179616	60835	53563	2,953	
4	reduced Nef-Tat	91726	73767	30948	20679	2,384	
5	adjuvant only	<4000	<4000	<4000	<4000		




Fig. 6B Tat-specific antibody titers and isotypes

midpoint titers						
group	immunization	Ig	IgG1	IgG2a	IgG2b	ratio IgG1/IgG2a
1	reduced Tat	212799	123242	62697	55763	1,966
2	reduced Nef-Tat	75676	84046	18449	11692	4,556
3	adjuvant only	<4000	<4000	<4000	<4000	




Fig. 7 Antigen-specific lymphoproliferative response of pooled lymph node cells

[3H] Thymidine incorporation in cpm				Data expressed as stimulation index			
	Group 1 reduced Tat	Group 2 reduced Nef-Tat	Group 3 adjuvant only		Group 1 reduced Tat	Group 2 reduced Nef-Tat	Group 3 adjuvant only
reduced Tat	41967	18511	789	reduced Tat	140	115	1
	37609	32346	415	5µg/ml	125	201	1
	27640	23408	397	1µg/ml	92	145	1
				0.2µg/ml			
reduced Nef-Tat	43882	31694	483	reduced Nef-Tat	146	197	1
	33865	28094	245	5µg/ml	113	174	0
	25079	22891	383	1µg/ml	84	142	1
	300	161	571	0.2µg/ml			
				medium	1	1	1








Fig. 8 Cell binding assay

Fig. 9 Inhibition of cell growth

SUBSTITUTE SHEET (RULE 26)

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: SmithKline Beecham Biologicals S.A.
- (ii) TITLE OF THE INVENTION: Vaccine
- (iii) NUMBER OF SEQUENCES: 27
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: SmithKline Beecham
 - (B) STREET: Two New Horizons Court
 - (C) CITY: Brentford
 - (D) STATE:
 - (E) COUNTRY: Middx, UK
 - (F) ZIP: TW8 9EP
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 26-SEP-1997
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Bor, Fiona R
 - (B) REGISTRATION NUMBER:
 - (C) REFERENCE/DOCKET NUMBER:
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 0181 975 2817
 - (B) TELEFAX: 0181 975 6141
 - (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATCGTCCATG .GGT.GGC.A AG.TGG.T

28

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CGGCTACTAG TGCAGTTCTT GAA

23

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATCGTACTAG T.GAG.CCA. GTA.GAT.C

29

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CGGCTACTAG TTCCTTCGG GCCT

24

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATCGTCCATG GAGCCAGTAG ATC

23

3 / 15

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 441 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

ATGGATCCAA AACTTTTAGC CCTTCTTTA TTAGCAGCTG GCGTACTAGC AGGTTGTAGC      60
AGCCATTCAT CAAATATGGC GAATACCCAA ATGAAATCAG ACAAATCAT TATTGCTCAC      120
CGTGGTGCTA GCGGTTATTT ACCAGAGCAT ACGTTAGAAT CTAAAGCACT TGCTTTTGCA      180
CAACAGGCTG ATTATTTAGA GCAAGATTTA GCAATGACTA AGGATGGTCG TTTAGTGGTT      240
ATTCACGATC ACTTTTTAGA TGGCTTGACT GATGTTGCGA AAAAATTCCC ACATCGTCAT      300
CGTAAAGATG GCCGTTACTA TGTATCGAC TTTACCTTAA AAGAAATTCA AAGTTTAGAA      360
ATGACAGAAA ACTTTGAAAC CATGGCCACG TGTGATCAGA GCTCACTAG TGGCCACCAT      420
CACCATCACC ATTAATCTAG A                                     441

```

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 144 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

```

Met Asp Pro Lys Thr Leu Ala Leu Ser Leu Leu Ala Ala Gly Val Leu
 1           5           10           15
Ala Gly Cys Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys
 20           25           30
Ser Asp Lys Ile Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro
 35           40           45
Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala Asp
 50           55           60
Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val
 65           70           75           80
Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe
 85           90           95
Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr
100          105          110
Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met
115          120          125
Ala Thr Cys Asp Gln Ser Ser Thr Ser Gly His His His His His His
130          135          140

```

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 648 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

SUBSTITUTE SHEET (RULE 26)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATGGGTGGCA	AGTGGTCAAA	AAGTAGTGTG	GTTGGATGGC	CTACTGTAAG	GGAAAGAATG	60
AGACGAGCTG	AGCCAGCAGC	AGATGGGGTG	GGAGCAGCAT	CTCGAGACCT	GGAAAAACAT	120
GGAGCAATCA	CAAGTAGCAA	TACAGCAGCT	ACCAATGCTG	CTTGTGCCTG	GCTAGAAGCA	180
CAAGAGGAGG	AGGAGGTGGG	TTTTCCAGTC	ACACCTCAGG	TACCTTTAAG	ACCAATGACT	240
TACAAGGCAG	CTGTAGATCT	TAGCCACTTT	TTAAAAGAAA	AGGGGGGACT	GGAAGGGCTA	300
ATTCACTCCC	AACGAAGACA	AGATATCCTT	GATCTGTGGA	TCTACCACAC	ACAAGGCTAC	360
TTCCCTGATT	GGCAGAACTA	CACACCAGGG	CCAGGGGTCA	GATATCCACT	GACCTTTGGA	420
TGGTGCTACA	AGCTAGTACC	AGTTGAGCCA	GATAAGGTAG	AAGAGGCCAA	TAAAGGAGAG	480
AACACCAGCT	TGTTACACCC	TGTGAGCCTG	CATGGAATGG	ATGACCCTGA	GAGAGAAGTG	540
TTAGAGTGGA	GGTTTGACAG	CCGCCTAGCA	TTTCATCACG	TGGCCCGAGA	GCTGCATCCG	600
GAGTACTTCA	AGAACTGCAC	TAGTGGCCAC	CATCACCATC	ACCATTA		648

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 216 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Met	Gly	Gly	Lys	Trp	Ser	Lys	Ser	Ser	Val	Val	Gly	Trp	Pro	Thr	Val	
1				5					10					15		
Arg	Glu	Arg	Met	Arg	Arg	Ala	Glu	Pro	Ala	Ala	Asp	Gly	Val	Gly	Ala	
			20					25					30			
Ala	Ser	Arg	Asp	Leu	Glu	Lys	His	Gly	Ala	Ile	Thr	Ser	Ser	Asn	Thr	
			35				40					45				
Ala	Ala	Thr	Asn	Ala	Ala	Cys	Ala	Trp	Leu	Glu	Ala	Gln	Glu	Glu	Glu	
			50			55					60					
Glu	Val	Gly	Phe	Pro	Val	Thr	Pro	Gln	Val	Pro	Leu	Arg	Pro	Met	Thr	
65					70					75					80	
Tyr	Lys	Ala	Ala	Val	Asp	Leu	Ser	His	Phe	Leu	Lys	Glu	Lys	Gly	Gly	
				85					90					95		
Leu	Glu	Gly	Leu	Ile	His	Ser	Gln	Arg	Arg	Gln	Asp	Ile	Leu	Asp	Leu	
			100					105					110			
Trp	Ile	Tyr	His	Thr	Gln	Gly	Tyr	Phe	Pro	Asp	Trp	Gln	Asn	Tyr	Thr	
		115				120						125				
Pro	Gly	Pro	Gly	Val	Arg	Tyr	Pro	Leu	Thr	Phe	Gly	Trp	Cys	Tyr	Lys	
		130				135					140					
Leu	Val	Pro	Val	Glu	Pro	Asp	Lys	Val	Glu	Glu	Ala	Asn	Lys	Gly	Glu	
145					150					155					160	
Asn	Thr	Ser	Leu	Leu	His	Pro	Val	Ser	Leu	His	Gly	Met	Asp	Asp	Pro	
			165						170					175		
Glu	Arg	Glu	Val	Leu	Glu	Trp	Arg	Phe	Asp	Ser	Arg	Leu	Ala	Phe	His	
			180					185					190			
His	Val	Ala	Arg	Glu	Leu	His	Pro	Glu	Tyr	Phe	Lys	Asn	Cys	Thr	Ser	
		195					200						205			
Gly	His	His	His	His	His	His										
		210				215										

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 288 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ATGGAGCCAG	TAGATCCTAG	ACTAGAGCCC	TGGAAGCATC	CAGGAAGTCA	GCCTAAAACT	60
GCTTGTTACCA	ATTGCTATTG	TAAAAAGTGT	TGCTTTTCATT	GCCAAGTTTG	TTTCATAACA	120
AAAGCCTTAG	GCATCTCCTA	TGGCAGGAAG	AAGCGGAGAC	AGCGACGAAG	ACCTCCTCAA	180
GGCAGTCAGA	CTCATCAAGT	TTCTCTATCA	AAGCAACCCA	CCTCCCAATC	CCGAGGGGAC	240
CCGACAGGCC	CGAAGGAAAC	TAGTGGCCAC	CATCACCATC	ACCATTAA		288

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 96 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met	Glu	Pro	Val	Asp	Pro	Arg	Leu	Glu	Pro	Trp	Lys	His	Pro	Gly	Ser
1				5				10					15		
Gln	Pro	Lys	Thr	Ala	Cys	Thr	Asn	Cys	Tyr	Cys	Lys	Lys	Cys	Cys	Phe
			20				25					30			
His	Cys	Gln	Val	Cys	Phe	Ile	Thr	Lys	Ala	Leu	Gly	Ile	Ser	Tyr	Gly
			35				40					45			
Arg	Lys	Lys	Arg	Arg	Gln	Arg	Arg	Pro	Pro	Gln	Gly	Ser	Gln	Thr	
			50			55				60					
His	Gln	Val	Ser	Leu	Ser	Lys	Gln	Pro	Thr	Ser	Gln	Ser	Arg	Gly	Asp
65				70					75						80
Pro	Thr	Gly	Pro	Lys	Glu	Thr	Ser	Gly	His	His	His	His	His	His	His
				85					90						95

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 909 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ATGGGTGGCA	AGTGGTCAA	AAGTAGTGTG	GTTGGATGGC	CTACTGTAAG	GGAAAGAATG	60
AGACGAGCTG	AGCCAGCAGC	AGATGGGGTG	GGAGCAGCAT	CTCGAGACCT	GGAAAAACAT	120
GGAGCAATCA	CAAGTAGCAA	TACAGCAGCT	ACCAATGCTG	CTTGTGCCTG	GCTAGAAGCA	180
CAAGAGGAGG	AGGAGGTGGG	TTTTCCAGTC	ACACCTCAGG	TACCTTTAAG	ACCAATGACT	240
TACAAGGCAG	CTGTAGATCT	TAGCCACTTT	TAAAAGAAA	AGGGGGGACT	GGAAGGGCTA	300
ATTCACTCCC	AACGAAGACA	AGATATCCTT	GATCTGTGGA	TCTACCACAC	ACAAGGCTAC	360

TTCCTGATT	GGCAGAACTA	CACACCAGGG	CCAGGGGTCA	GATATCCACT	GACCTTTGGA	420
TGGTGCTACA	AGCTAGTACC	AGTTGAGCCA	GATAAGGTAG	AAGAGGCCAA	TAAAGGAGAG	480
AACACCAGCT	TGTTACACCC	TGTGAGCCTG	CATGGAATGG	ATGACCCTGA	GAGAGAAGTG	540
TTAGAGTGGA	GGTTTGACAG	CCGCCTAGCA	TTTCATCACG	TGGCCCGAGA	GCTGCATCCG	600
GAGTACTTCA	AGAACTGCAC	TAGTGAGCCA	GTAGATCCTA	GACTAGAGCC	CTGGAAGCAT	660
CCAGGAAGTC	AGCCTAAAAC	TGCTTGTAAC	AATTGCTATT	GTAAAAAGTG	TTGCTTTCAT	720
TGCCAAGTTT	GTTTCATAAC	AAAAGCCTTA	GGCATCTCCT	ATGGCAGGAA	GAAGCGGAGA	780
CAGCGACGAA	GACCTCCTCA	AGGCAGTCAG	ACTCATCAAG	TTTCTCTATC	AAAGCAACCC	840
ACCTCCCAAT	CCCGAGGGGA	CCCACAGGC	CCGAAGGAAA	CTAGTGGCCA	CCATCACCAT	900
CACCATTAA						909

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 303 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Met	Gly	Gly	Lys	Trp	Ser	Lys	Ser	Ser	Val	Val	Gly	Trp	Pro	Thr	Val
1				5					10					15	
Arg	Glu	Arg	Met	Arg	Arg	Ala	Glu	Pro	Ala	Ala	Asp	Gly	Val	Gly	Ala
			20					25					30		
Ala	Ser	Arg	Asp	Leu	Glu	Lys	His	Gly	Ala	Ile	Thr	Ser	Ser	Asn	Thr
		35				40						45			
Ala	Ala	Thr	Asn	Ala	Ala	Cys	Ala	Trp	Leu	Glu	Ala	Gln	Glu	Glu	Glu
	50					55					60				
Glu	Val	Gly	Phe	Pro	Val	Thr	Pro	Gln	Val	Pro	Leu	Arg	Pro	Met	Thr
65					70				75					80	
Tyr	Lys	Ala	Ala	Val	Asp	Leu	Ser	His	Phe	Leu	Lys	Glu	Lys	Gly	Gly
			85						90					95	
Leu	Glu	Gly	Leu	Ile	His	Ser	Gln	Arg	Arg	Gln	Asp	Ile	Leu	Asp	Leu
			100					105					110		
Trp	Ile	Tyr	His	Thr	Gln	Gly	Tyr	Phe	Pro	Asp	Trp	Gln	Asn	Tyr	Thr
		115				120						125			
Pro	Gly	Pro	Gly	Val	Arg	Tyr	Pro	Leu	Thr	Phe	Gly	Trp	Cys	Tyr	Lys
		130				135					140				
Leu	Val	Pro	Val	Glu	Pro	Asp	Lys	Val	Glu	Glu	Ala	Asn	Lys	Gly	Glu
145					150					155				160	
Asn	Thr	Ser	Leu	Leu	His	Pro	Val	Ser	Leu	His	Gly	Met	Asp	Asp	Pro
			165					170						175	
Glu	Arg	Glu	Val	Leu	Glu	Trp	Arg	Phe	Asp	Ser	Arg	Leu	Ala	Phe	His
			180					185					190		
His	Val	Ala	Arg	Glu	Leu	His	Pro	Glu	Tyr	Phe	Lys	Asn	Cys	Thr	Ser
		195					200					205			
Glu	Pro	Val	Asp	Pro	Arg	Leu	Glu	Pro	Trp	Lys	His	Pro	Gly	Ser	Gln
		210				215					220				
Pro	Lys	Thr	Ala	Cys	Thr	Asn	Cys	Tyr	Cys	Lys	Lys	Cys	Cys	Phe	His
225					230					235				240	
Cys	Gln	Val	Cys	Phe	Ile	Thr	Lys	Ala	Leu	Gly	Ile	Ser	Tyr	Gly	Arg
			245						250					255	
Lys	Lys	Arg	Arg	Gln	Arg	Arg	Arg	Pro	Pro	Gln	Gly	Ser	Gln	Thr	His
			260					265					270		
Gln	Val	Ser	Leu	Ser	Lys	Gln	Pro	Thr	Ser	Gln	Ser	Arg	Gly	Asp	Pro

Thr Gly Pro Lys Glu Thr Ser Gly His His His His His His
275 280 285
290 295 300

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- ```
(A) LENGTH: 1029 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

|             |            |            |            |            |            |      |
|-------------|------------|------------|------------|------------|------------|------|
| ATGGATCCAA  | AAACTTTAGC | CCTTTCTTTA | TTAGCAGCTG | GCGTACTAGC | AGGTTGTAGC | 60   |
| AGCCATTCAT  | CAAATATGGC | GAATACCCAA | ATGAAATCAG | ACAAAATCAT | TATTGCTCAC | 120  |
| CGTGGTGCTA  | GCGGTTATTT | ACCAGAGCAT | ACGTTAGAAT | CTAAAGCACT | TGCTTTTGCA | 180  |
| CAACAGGCTG  | ATTATTTAGA | GCAAGATTTA | GCAATGACTA | AGGATGGTCG | TTTAGTGGTT | 240  |
| ATTCACGATC  | ACCTTTTAGA | TGGCTTGACT | GATGTTGCGA | AAAAATTCCC | ACATCGTCAT | 300  |
| CGTAAAGATG  | GCCGTTACTA | TGTCATCGAC | TTTACCTTAA | AAGAAATTCA | AAGTTTAGAA | 360  |
| ATGACAGAAA  | ACTTTGAAAC | CATGGGTGGC | AAGTGGTCAA | AAAGTAGTGT | GGTTGGATGG | 420  |
| CCTACTGTAA  | GGGAAAAGAT | GAGACGAGCT | GAGCCAGCAG | CAGATGGGGT | GGGAGCAGCA | 480  |
| TCTCGAGACC  | TGGAAAAACA | TGGAGCAATC | ACAAGTAGCA | ATACAGCAGC | TACCAATGCT | 540  |
| GCTTGTGCCT  | GGCTAGAAGC | ACAAGAGGAG | GAGGAGGTGG | GTTTTCCAGT | CACACCTCAG | 600  |
| GTACCTTTAA  | GACCAATGAC | TTACAAGGCA | GCTGTAGATC | TTAGCCACTT | TTTAAAAGAA | 660  |
| AAGGGGGGAC  | TGGAAGGGCT | AATTCACTCC | CAACGAAGAC | AAGATATCCT | TGATCTGTGG | 720  |
| ATCTACCACA  | CACAAGGCTA | CTTCCCTGAT | TGGCAGAACT | ACACACCAGG | GCCAGGGGTC | 780  |
| AGATATCCAC  | TGACCTTTGG | ATGGTGCTAC | AAGCTAGTAC | CAGTTGAGCC | AGATAAGGTA | 840  |
| GAAGAGGCCA  | ATAAAGGAGA | GAACACCAGC | TGTGTACACC | CTGTGAGCCT | GCATGGAATG | 900  |
| GATGACCCTG  | AGAGAGAAGT | GGTAGAGTGG | AGGTTTGACA | GCCGCCTAGC | ATTTCATCAC | 960  |
| CTGGCCCCGAG | AGCTGCATCC | GGAGTACTTC | AAGAAGTGCA | CTAGTGGCCA | CCATCACCAT | 1020 |
| CACCATTA    |            |            |            |            |            | 1029 |

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 325 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

|           |           |           |     |           |           |           |           |           |           |           |           |           |           |           |     |
|-----------|-----------|-----------|-----|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----|
| Cys<br>1  | Ser       | Ser       | His | Ser<br>5  | Ser       | Asn       | Met       | Ala       | Asn<br>10 | Thr       | Gln       | Met       | Lys       | Ser<br>15 | Asp |
| Lys       | Ile       | Ile       | Ile | Ala<br>20 | His       | Arg       | Gly       | Ala<br>25 | Ser       | Gly       | Tyr       | Leu       | Pro<br>30 | Glu       | His |
| Thr       | Leu       | Glu<br>35 | Ser | Lys       | Ala       | Leu       | Ala<br>40 | Phe       | Ala       | Gln       | Gln       | Ala<br>45 | Asp       | Tyr       | Leu |
| Glu       | Gln<br>50 | Asp       | Leu | Ala       | Met       | Thr<br>55 | Lys       | Asp       | Gly       | Arg       | Leu<br>60 | Val       | Val       | Ile       | His |
| Asp<br>65 | His       | Phe       | Leu | Asp       | Gly<br>70 | Leu       | Thr       | Asp       | Val       | Ala<br>75 | Lys       | Lys       | Phe       | Pro<br>80 | His |
| Arg       | His       | Arg       | Lys | Asp<br>85 | Gly       | Arg       | Tyr       | Tyr       | Val<br>90 | Ile       | Asp       | Phe       | Thr       | Leu<br>95 | Lys |

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |  |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| Glu | Ile | Gln | Ser | Leu | Glu | Met | Thr | Glu | Asn | Phe | Glu | Thr | Met | Gly | Gly |  |  |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     | 110 |     |     |  |  |
| Lys | Trp | Ser | Lys | Ser | Ser | Val | Val | Gly | Trp | Pro | Thr | Val | Arg | Glu | Arg |  |  |
|     |     | 115 |     |     |     |     | 120 |     |     |     |     | 125 |     |     |     |  |  |
| Met | Arg | Arg | Ala | Glu | Pro | Ala | Ala | Asp | Gly | Val | Gly | Ala | Ala | Ser | Arg |  |  |
|     | 130 |     |     |     |     | 135 |     |     |     |     | 140 |     |     |     |     |  |  |
| Asp | Leu | Glu | Lys | His | Gly | Ala | Ile | Thr | Ser | Ser | Asn | Thr | Ala | Ala | Thr |  |  |
| 145 |     |     |     |     | 150 |     |     |     |     | 155 |     |     |     |     | 160 |  |  |
| Asn | Ala | Ala | Cys | Ala | Trp | Leu | Glu | Ala | Gln | Glu | Glu | Glu | Glu | Val | Gly |  |  |
|     |     |     | 165 |     |     |     |     |     | 170 |     |     |     |     | 175 |     |  |  |
| Phe | Pro | Val | Thr | Pro | Gln | Val | Pro | Leu | Arg | Pro | Met | Thr | Tyr | Lys | Ala |  |  |
|     |     |     | 180 |     |     |     |     | 185 |     |     |     |     |     | 190 |     |  |  |
| Ala | Val | Asp | Leu | Ser | His | Phe | Leu | Lys | Glu | Lys | Gly | Gly | Leu | Glu | Gly |  |  |
|     | 195 |     |     |     |     |     | 200 |     |     |     |     | 205 |     |     |     |  |  |
| Leu | Ile | His | Ser | Gln | Arg | Arg | Gln | Asp | Ile | Leu | Asp | Leu | Trp | Ile | Tyr |  |  |
|     | 210 |     |     |     |     | 215 |     |     |     |     | 220 |     |     |     |     |  |  |
| His | Thr | Gln | Gly | Tyr | Phe | Pro | Asp | Trp | Gln | Asn | Tyr | Thr | Pro | Gly | Pro |  |  |
| 225 |     |     |     |     | 230 |     |     |     |     | 235 |     |     |     |     | 240 |  |  |
| Gly | Val | Arg | Tyr | Pro | Leu | Thr | Phe | Gly | Trp | Cys | Tyr | Lys | Leu | Val | Pro |  |  |
|     |     |     | 245 |     |     |     |     | 250 |     |     |     |     |     | 255 |     |  |  |
| Val | Glu | Pro | Asp | Lys | Val | Glu | Glu | Ala | Asn | Lys | Gly | Glu | Asn | Thr | Ser |  |  |
|     |     |     | 260 |     |     |     |     | 265 |     |     |     |     | 270 |     |     |  |  |
| Leu | Leu | His | Pro | Val | Ser | Leu | His | Gly | Met | Asp | Asp | Pro | Glu | Arg | Glu |  |  |
|     |     | 275 |     |     |     |     | 280 |     |     |     |     | 285 |     |     |     |  |  |
| Val | Leu | Glu | Trp | Arg | Phe | Asp | Ser | Arg | Leu | Ala | Phe | His | His | Val | Ala |  |  |
|     | 290 |     |     |     |     | 295 |     |     |     |     | 300 |     |     |     |     |  |  |
| Arg | Glu | Leu | His | Pro | Glu | Tyr | Phe | Lys | Asn | Cys | Thr | Ser | Gly | His | His |  |  |
| 305 |     |     |     |     | 310 |     |     |     | 315 |     |     |     |     |     | 320 |  |  |
| His | His | His | His |     |     |     |     |     |     |     |     |     |     |     |     |  |  |

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1290 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

|            |            |            |            |            |            |     |
|------------|------------|------------|------------|------------|------------|-----|
| ATGGATCCAA | AAACTTTAGC | CCTTTCTTTA | TTAGCAGCTG | GCGTACTAGC | AGGTTGTAGC | 60  |
| AGCCATTCAT | CAAATATGGC | GAATACCCAA | ATGAAATCAG | ACAAAATCAT | TATTGCTCAC | 120 |
| CGTGGTGCTA | GCGGTTATTT | ACCAGAGCAT | ACGTTAGAAT | CTAAAGCACT | TGCGTTTGCA | 180 |
| CAACAGGCTG | ATTATTTAGA | GCAAGATTTA | GCAATGACTA | AGGATGGTCG | TTTAGTGGTT | 240 |
| ATTCACGATC | ACTTTTTAGA | TGGCTTGACT | GATGTTGCGA | AAAAATTCCC | ACATCGTCAT | 300 |
| CGTAAAGATG | GCCGTTACTA | TGTCATCGAC | TTTACCTTAA | AAGAAATTCA | AAGTTTAGAA | 360 |
| ATGACAGAAA | ACTTTGAAAC | CATGGGTGGC | AAGTGGTCAA | AAAGTAGTGT | GGTTGGATGG | 420 |
| CCTACTGTAA | GGGAAAGAAT | GAGACGAGCT | GAGCCAGCAG | CAGATGGGGT | GGGAGCAGCA | 480 |
| TCTCGAGACC | TGGAAAAACA | TGGAGCAATC | ACAAGTAGCA | ATACAGCAGC | TACCAATGCT | 540 |
| GCTTGTGCCT | GGCTAGAAGC | ACAAGAGGAG | GAGGAGGTGG | GTTTTCCAGT | CACACCTCAG | 600 |
| GTACCTTTAA | GACCAATGAC | TTACAAGGCA | GCTGTAGATC | TTAGCCACTT | TTTAAAAGAA | 660 |
| AAGGGGGGAC | TGGAAGGGCT | AATTCACCTC | CAACGAAGAC | AAGATATCCT | TGATCTGTGG | 720 |
| ATCTACCACA | CACAAGGCTA | CTTCCCTGAT | TGGCAGAACT | ACACACCAGG | GCCAGGGGTC | 780 |
| AGATATCCAC | TGACCTTTGG | ATGGTGCTAC | AAGCTAGTAC | CAGTTGAGCC | AGATAAGGTA | 840 |
| GAAGAGGCCA | ATAAAGGAGA | GAACACCAGC | TTGTTACACC | CTGTGAGCCT | GCATGGAATG | 900 |



|            |            |            |            |            |             |      |
|------------|------------|------------|------------|------------|-------------|------|
| GATGACCCTG | AGAGAGAAGT | GTTAGAGTGG | AGGTTTGACA | GCCGCCTAGC | ATTTTCATCAC | 960  |
| GTGGCCCGAG | AGCTGCATCC | GGAGTACTTC | AAGAACTGCA | CTAGTGAGCC | AGTAGATCCT  | 1020 |
| AGACTAGAGC | CCTGGAAGCA | TCCAGGAAGT | CAGCCTAAAA | CTGCTTGAC  | CAATTGCTAT  | 1080 |
| TGTAAAAAGT | GTTGCTTTCA | TTGCCAAGTT | TGTTTCATAA | CAAAAGCCTT | AGGCATCTCC  | 1140 |
| TATGGCAGGA | AGAAGCGGAG | ACAGCGACGA | AGACCTCCTC | AAGGCAGTCA | GACTCATCAA  | 1200 |
| GTTTCTCTAT | CAAAGCAACC | CACCTCCCAA | TCCCGAGGGG | ACCCGACAGG | CCCGAAGGAA  | 1260 |
| ACTAGTGGCC | ACCATCACCA | TCACCATTAA |            |            |             | 1290 |

## (2) INFORMATION FOR SEQ ID NO:17:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 412 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Cys | Ser | Ser | His | Ser | Ser | Asn | Met | Ala | Asn | Thr | Gln | Met | Lys | Ser | Asp | 1   | 5   | 10  | 15  |
| Lys | Ile | Ile | Ile | Ala | His | Arg | Gly | Ala | Ser | Gly | Tyr | Leu | Pro | Glu | His | 20  | 25  | 30  |     |
| Thr | Leu | Glu | Ser | Lys | Ala | Leu | Ala | Phe | Ala | Gln | Gln | Ala | Asp | Tyr | Leu | 35  | 40  | 45  |     |
| Glu | Gln | Asp | Leu | Ala | Met | Thr | Lys | Asp | Gly | Arg | Leu | Val | Val | Ile | His | 50  | 55  | 60  |     |
| Asp | His | Phe | Leu | Asp | Gly | Leu | Thr | Asp | Val | Ala | Lys | Lys | Phe | Pro | His | 65  | 70  | 75  | 80  |
| Arg | His | Arg | Lys | Asp | Gly | Arg | Tyr | Tyr | Val | Ile | Asp | Phe | Thr | Leu | Lys | 85  | 90  | 95  |     |
| Glu | Ile | Gln | Ser | Leu | Glu | Met | Thr | Glu | Asn | Phe | Glu | Thr | Met | Gly | Gly | 100 | 105 | 110 |     |
| Lys | Trp | Ser | Lys | Ser | Ser | Val | Val | Gly | Trp | Pro | Thr | Val | Arg | Glu | Arg | 115 | 120 | 125 |     |
| Met | Arg | Arg | Ala | Glu | Pro | Ala | Ala | Asp | Gly | Val | Gly | Ala | Ala | Ser | Arg | 130 | 135 | 140 |     |
| Asp | Leu | Glu | Lys | His | Gly | Ala | Ile | Thr | Ser | Ser | Asn | Thr | Ala | Ala | Thr | 145 | 150 | 155 | 160 |
| Asn | Ala | Ala | Cys | Ala | Trp | Leu | Glu | Ala | Gln | Glu | Glu | Glu | Glu | Val | Gly | 165 | 170 | 175 |     |
| Phe | Pro | Val | Thr | Pro | Gln | Val | Pro | Leu | Arg | Pro | Met | Thr | Tyr | Lys | Ala | 180 | 185 | 190 |     |
| Ala | Val | Asp | Leu | Ser | His | Phe | Leu | Lys | Glu | Lys | Gly | Gly | Leu | Glu | Gly | 195 | 200 | 205 |     |
| Leu | Ile | His | Ser | Gln | Arg | Arg | Gln | Asp | Ile | Leu | Asp | Leu | Trp | Ile | Tyr | 210 | 215 | 220 |     |
| His | Thr | Gln | Gly | Tyr | Phe | Pro | Asp | Trp | Gln | Asn | Tyr | Thr | Pro | Gly | Pro | 225 | 230 | 235 | 240 |
| Gly | Val | Arg | Tyr | Pro | Leu | Thr | Phe | Gly | Trp | Cys | Tyr | Lys | Leu | Val | Pro | 245 | 250 | 255 |     |
| Val | Glu | Pro | Asp | Lys | Val | Glu | Glu | Ala | Asn | Lys | Gly | Glu | Asn | Thr | Ser | 260 | 265 | 270 |     |
| Leu | Leu | His | Pro | Val | Ser | Leu | His | Gly | Met | Asp | Asp | Pro | Glu | Arg | Glu | 275 | 280 | 285 |     |
| Val | Leu | Glu | Trp | Arg | Phe | Asp | Ser | Arg | Leu | Ala | Phe | His | His | Val | Ala | 290 | 295 | 300 |     |

10 / 15

```

Arg Glu Leu His Pro Glu Tyr Phe Lys Asn Cys Thr Ser Glu Pro Val
305 310 315 320
Asp Pro Arg Leu Glu Pro Trp Lys His Pro Gly Ser Gln Pro Lys Thr
 325 330 335
Ala Cys Thr Asn Cys Tyr Cys Lys Lys Cys Cys Phe His Cys Gln Val
 340 345 350
Cys Phe Ile Thr Lys Ala Leu Gly Ile Ser Tyr Gly Arg Lys Lys Arg
 355 360 365
Arg Gln Arg Arg Arg Pro Pro Gln Gly Ser Gln Thr His Gln Val Ser
 370 375 380
Leu Ser Lys Gln Pro Thr Ser Gln Ser Arg Gly Asp Pro Thr Gly Pro
385 390 395 400
Lys Glu Thr Ser Gly His His His His His His
 405 410

```

## (2) INFORMATION FOR SEQ ID NO:18:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 981 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

```

ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC 60
ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA 120
CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT 180
CGTTTAGTGG TTATTCACGA TCACTTTTTT GATGGCTTGA CTGATGTTGC GAAAAAATTC 240
CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATT 300
CAAAGTTTAG AAATGACAGA AACTTTTGAA ACCATGGGTG GCAAGTGGTC AAAAAGTAGT 360
GTGGTTGGAT GGCCTACTGT AAGGGAAAGA ATGAGACGAG CTGAGCCAGC AGCAGATGGG 420
GTGGGAGCAG CATCTCGAGA CCTGGAAAAA CATGGAGCAA TCACAAGTAG CAATACAGCA 480
GCTACCAATG CTGCTTGTGC CTGGCTAGAA GCACAAGAGG AGGAGGAGGT GGGTTTTCCA 540
GTCACACCTC AGGTACCTTT AAGACCAATG ACTTACAAGG CAGCTGTAGA TCTTAGCCAC 600
TTTTTAAAG AAAAGGGGGG ACTGGAAGGG CTAATTCAC CCCAACGAAG ACAAGATATC 660
CTTGATCTGT GGATCTACCA CACACAAGGC TACTTCCCTG ATTGGCAGAA CTACACACCA 720
GGGCCAGGGG TCAGATATCC ACTGACCTTT GGATGGTGCT ACAAGCTAGT ACCAGTTGAG 780
CCAGATAAGG TAGAAGAGGC CAATAAAGGA GAGAACACCA GCTTGTTACA CCCTGTGAGC 840
CTGCATGGAA TGGATGACCC TGAGAGAGAA GTGTTAGAGT GGAGGTTTGA CAGCCGCCTA 900
GCATTCATC ACGTGGCCCG AGAGCTGCAT CCGGAGTACT TCAAGAACTG CACTAGTGGC 960
CACCATCACC ATCACCATTA A

```

## (2) INFORMATION FOR SEQ ID NO:19:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 327 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```

Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys
1 5 10 15

```

```

Ser Asp Lys Ile Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro
 20 25 30
Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala Asp
 35 40 45
Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val
 50 55 60
Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe
 65 70 75 80
Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr
 85 90 95
Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met
 100 105 110
Gly Gly Lys Trp Ser Lys Ser Ser Val Val Gly Trp Pro Thr Val Arg
 115 120 125
Glu Arg Met Arg Arg Ala Glu Pro Ala Ala Asp Gly Val Gly Ala Ala
 130 135 140
Ser Arg Asp Leu Glu Lys His Gly Ala Ile Thr Ser Ser Asn Thr Ala
 145 150 155 160
Ala Thr Asn Ala Ala Cys Ala Trp Leu Glu Ala Gln Glu Glu Glu Glu
 165 170 175
Val Gly Phe Pro Val Thr Pro Gln Val Pro Leu Arg Pro Met Thr Tyr
 180 185 190
Lys Ala Ala Val Asp Leu Ser His Phe Leu Lys Glu Lys Gly Gly Leu
 195 200 205
Glu Gly Leu Ile His Ser Gln Arg Arg Gln Asp Ile Leu Asp Leu Trp
 210 215 220
Ile Tyr His Thr Gln Gly Tyr Phe Pro Asp Trp Gln Asn Tyr Thr Pro
 225 230 235 240
Gly Pro Gly Val Arg Tyr Pro Leu Thr Phe Gly Trp Cys Tyr Lys Leu
 245 250 255
Val Pro Val Glu Pro Asp Lys Val Glu Glu Ala Asn Lys Gly Glu Asn
 260 265 270
Thr Ser Leu Leu His Pro Val Ser Leu His Gly Met Asp Asp Pro Glu
 275 280 285
Arg Glu Val Leu Glu Trp Arg Phe Asp Ser Arg Leu Ala Phe His His
 290 295 300
Val Ala Arg Glu Leu His Pro Glu Tyr Phe Lys Asn Cys Thr Ser Gly
 305 310 315 320
His His His His His His
 325

```

## (2) INFORMATION FOR SEQ ID NO:20:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1242 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

```

ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC 60
ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA 120
CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT 180
CGTTTAGTGG TTATTCACGA TCACTTTTGA GATGGCTTGA CTGATGTTGC GAAAAAATTC 240
CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATT 300

```

```

CAAAGTTTAG AAATGACAGA AACTTTTGAA ACCATGGGTG GCAAGTGGTC AAAAAGTAGT 360
GTGGTTGGAT GGCCTACTGT AAGGGAAAGA ATGAGACGAG CTGAGCCAGC AGCAGATGGG 420
GTGGGAGCAG CATCTCGAGA CCTGGAAAAA CATGGAGCAA TCACAAGTAG CAATACAGCA 480
GCTACCAATG CTGCTTGTGC CTGGCTAGAA GCACAAGAGG AGGAGGAGGT GGGTTTTCCA 540
GTCACACCTC AGGTACCTTT AAGACCAATG ACTTACAAGG CAGCTGTAGA TCTTAGCCAC 600
TTTTTAAAAG AAAAGGGGGG ACTGGAAGGG CTAATTCCT CCCAACGAAG ACAAGATATC 660
CTTGATCTGT GGATCTACCA CACACAAGGC TACTTCCCTG ATTGGCAGAA CTACACACCA 720
GGGCCAGGGG TCAGATATCC ACTGACCTTT GGATGGTGCT ACAAGCTAGT ACCAGTTGAG 780
CCAGATAAGG TAGAAGAGGC CAATAAAGGA GAGAACACCA GCTTGTTACA CCCTGTGAGC 840
CTGCATGGAA TGGATGACCC TGAGAGAGAA GTGTTAGAGT GGAGGTTTGA CAGCCGCCTA 900
GCATTTTCATC ACGTGGCCCG AGAGCTGCAT CCGGAGTACT TCAAGAACTG CACTAGTGAG 960
CCAGTAGATC CTAGACTAGA GCCCTGGAAG CATCCAGGAA GTCAGCCTAA AACTGCTTGT 1020
ACCAATTGCT ATTGTAAAAA GTGTTGCTTT CATTGCCAAG TTTGTTTCAT AACAAAAGCC 1080
TTAGGCATCT CCTATGGCAG GAAGAAGCGG AGACAGCGAC GAAGACCTCC TCAAGGCAGT 1140
CAGACTCATC AAGTTTCTCT ATCAAAGCAA CCCACCTCCC AATCCCGAGG GGACCCGACA 1200
GGCCCGAAGG AACTAGTGG CCACCATCAC CATCACCATT AA 1242

```

## (2) INFORMATION FOR SEQ ID NO:21:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 414 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```

Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys
1 5 10 15
Ser Asp Lys Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro
20 25 30
Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala Asp
35 40 45
Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val
50 55 60
Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe
65 70 75 80
Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr
85 90 95
Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met
100 105 110
Gly Gly Lys Trp Ser Lys Ser Ser Val Val Gly Trp Pro Thr Val Arg
115 120 125
Glu Arg Met Arg Arg Ala Glu Pro Ala Ala Asp Gly Val Gly Ala Ala
130 135 140
Ser Arg Asp Leu Glu Lys His Gly Ala Ile Thr Ser Ser Asn Thr Ala
145 150 155 160
Ala Thr Asn Ala Ala Cys Ala Trp Leu Glu Ala Gln Glu Glu Glu
165 170 175
Val Gly Phe Pro Val Thr Pro Gln Val Pro Leu Arg Pro Met Thr Tyr
180 185 190
Lys Ala Ala Val Asp Leu Ser His Phe Leu Lys Glu Lys Gly Gly Leu
195 200 205
Glu Gly Leu Ile His Ser Gln Arg Arg Gln Asp Ile Leu Asp Leu Trp
210 215 220
Ile Tyr His Thr Gln Gly Tyr Phe Pro Asp Trp Gln Asn Tyr Thr Pro

```

13 / 15

```

225 230 235 240
Gly Pro Gly Val Arg Tyr Pro Leu Thr Phe Gly Trp Cys Tyr Lys Leu
245 250 255
Val Pro Val Glu Pro Asp Lys Val Glu Glu Ala Asn Lys Gly Glu Asn
260 265 270
Thr Ser Leu Leu His Pro Val Ser Leu His Gly Met Asp Asp Pro Glu
275 280 285
Arg Glu Val Leu Glu Trp Arg Phe Asp Ser Arg Leu Ala Phe His His
290 295 300
Val Ala Arg Glu Leu His Pro Glu Tyr Phe Lys Asn Cys Thr Ser Glu
305 310 315 320
Pro Val Asp Pro Arg Leu Glu Pro Trp Lys His Pro Gly Ser Gln Pro
325 330 335
Lys Thr Ala Cys Thr Asn Cys Tyr Cys Lys Lys Cys Cys Phe His Cys
340 345 350
Gln Val Cys Phe Ile Thr Lys Ala Leu Gly Ile Ser Tyr Gly Arg Lys
355 360 365
Lys Arg Arg Gln Arg Arg Arg Pro Pro Gln Gly Ser Gln Thr His Gln
370 375 380
Val Ser Leu Ser Lys Gln Pro Thr Ser Gln Ser Arg Gly Asp Pro Thr
385 390 395 400
Gly Pro Lys Glu Thr Ser Gly His His His His His His
405 410

```

## (2) INFORMATION FOR SEQ ID NO:22:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 288 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

ATGGAGCCAG TAGATCCTAG ACTAGAGCCC TGGAAGCATC CAGGAAGTCA GCCTAAACT 60
GCTTGTAACA ATTGCTATTG TAAAAAGTGT TGCTTTCATT GCCAAGTTTG TTTCATAACA 120
GCTGCCTTAG GCATCTCCTA TGGCAGGAAG AAGCGGAGAC AGCGACGAAG ACCTCCTCAA 180
GGCAGTCAGA CTCATCAAGT TTCTCTATCA AAGCAACCCA CCTCCCAATC CAAAGGGGAG 240
CCGACAGGCC CGAAGGAAAC TAGTGGCCAC CATCACCATC ACCATTAA 288

```

## (2) INFORMATION FOR SEQ ID NO:23:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 96 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

```

Met Glu Pro Val Asp Pro Arg Leu Glu Pro Trp Lys His Pro Gly Ser
1 5 10 15
Gln Pro Lys Thr Ala Cys Thr Asn Cys Tyr Cys Lys Lys Cys Cys Phe
20 25 30
His Cys Gln Val Cys Phe Ile Thr Ala Ala Leu Gly Ile Ser Tyr Gly

```

14 / 15

```

 35 40 45
Arg Lys Lys Arg Arg Gln Arg Arg Arg Pro Pro Gln Gly Ser Gln Thr
 50 55 60
His Gln Val Ser Leu Ser Lys Gln Pro Thr Ser Gln Ser Lys Gly Glu
 65 70 75 80
Pro Thr Gly Pro Lys Glu Thr Ser Gly His His His His His His
 85 90 95

```

## (2) INFORMATION FOR SEQ ID NO:24:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 909 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

```

ATGGGTGGCA AGTGGTCAAA AAGTAGTGTG GTTGGATGGC CTACTGTAAG GGAAAGAATG 60
AGACGAGCTG AGCCAGCAGC AGATGGGGTG GGAGCAGCAT CTCGAGACCT GGAAAAACAT 120
GGAGCAATCA CAAGTAGCAA TACAGCAGCT ACCAATGCTG CTTGTGCCTG GCTAGAAGCA 180
CAAGAGGAGG AGGAGGTGGG TTTTCCAGTC ACACCTCAGG TACCTTTAAG ACCAATGACT 240
TACAAGGCAG CTGTAGATCT TAGCCACTTT TTAAAAGAAA AGGGGGGACT GGAAGGGCTA 300
ATTCACCTCC AACGAAGACA AGATATCCTT GATCTGTGGA TCTACCACAC ACAAGGCTAC 360
TTCCCTGATT GGCAGAACTA CACACCAGGG CCAGGGGTCA GATATCCACT GACCTTTGGA 420
TGGTGCTACA AGCTAGTACC AGTTGAGCCA GATAAGGTAG AAGAGGCCAA TAAAGGAGAG 480
AACACCAGCT TGTTACACCC TGTGAGCCTG CATGGAATGG ATGACCCTGA GAGAGAAGTG 540
TTAGAGTGGA GGTTCGACAG CCGCCTAGCA TTTCATCAGC TGGCCCGAGA GCTGCATCCG 600
GAGTACTTCA AGAACTGCAC TAGTGAGCCA GTAGATCCTA GACTAGAGCC CTGGAAGCAT 660
CCAGGAAGTC AGCCTAAAAC TGCTTGTAAC AATTGCTATT GTAAAAAGTG TTGCTTTCAT 720
TGCCAAGTTT GTTTCATAAC AGCTGCCTTA GGCATCTCCT ATGGCAGGAA GAAGCGGAGA 780
CAGCGACGAA GACCTCCTCA AGGCAGTCAG ACTCATCAAG TTTCTCTATC AAAGCAACCC 840
ACCTCCCAAT CCAAAGGGGA GCCGACAGGC CCGAAGGAAA CTAGTGGCCA CCATCACCAT 900
CACCATTAA

```

## (2) INFORMATION FOR SEQ ID NO:25:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 303 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

```

Met Gly Gly Lys Trp Ser Lys Ser Ser Val Val Gly Trp Pro Thr Val
 1 5 10 15
Arg Glu Arg Met Arg Arg Ala Glu Pro Ala Ala Asp Gly Val Gly Ala
 20 25 30
Ala Ser Arg Asp Leu Glu Lys His Gly Ala Ile Thr Ser Ser Asn Thr
 35 40 45
Ala Ala Thr Asn Ala Ala Cys Ala Trp Leu Glu Ala Gln Glu Glu Glu
 50 55 60
Glu Val Gly Phe Pro Val Thr Pro Gln Val Pro Leu Arg Pro Met Thr
 65 70 75 80

```

SUBSTITUTE SHEET (RULE 26)

15 / 15

Tyr Lys Ala Ala Val Asp Leu Ser His Phe Leu Lys Glu Lys Gly Gly  
 85 90 95  
 Leu Glu Gly Leu Ile His Ser Gln Arg Arg Gln Asp Ile Leu Asp Leu  
 100 105 110  
 Trp Ile Tyr His Thr Gln Gly Tyr Phe Pro Asp Trp Gln Asn Tyr Thr  
 115 120 125  
 Pro Gly Pro Gly Val Arg Tyr Pro Leu Thr Phe Gly Trp Cys Tyr Lys  
 130 135 140  
 Leu Val Pro Val Glu Pro Asp Lys Val Glu Glu Ala Asn Lys Gly Glu  
 145 150 155 160  
 Asn Thr Ser Leu Leu His Pro Val Ser Leu His Gly Met Asp Asp Pro  
 165 170 175  
 Glu Arg Glu Val Leu Glu Trp Arg Phe Asp Ser Arg Leu Ala Phe His  
 180 185 190  
 His Val Ala Arg Glu Leu His Pro Glu Tyr Phe Lys Asn Cys Thr Ser  
 195 200 205  
 Glu Pro Val Asp Pro Arg Leu Glu Pro Trp Lys His Pro Gly Ser Gln  
 210 215 220  
 Pro Lys Thr Ala Cys Thr Asn Cys Tyr Cys Lys Lys Cys Cys Phe His  
 225 230 235 240  
 Cys Gln Val Cys Phe Ile Thr Ala Ala Leu Gly Ile Ser Tyr Gly Arg  
 245 250 255  
 Lys Lys Arg Arg Gln Arg Arg Arg Pro Pro Gln Gly Ser Gln Thr His  
 260 265 270  
 Gln Val Ser Leu Ser Lys Gln Pro Thr Ser Gln Ser Lys Gly Glu Pro  
 275 280 285  
 Thr Gly Pro Lys Glu Thr Ser Gly His His His His His His  
 290 295 300

## (2) INFORMATION FOR SEQ ID NO:26:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

TTCGAAACCA TGGCCGCGGA CTAGTGGCCA CCATCACCAT CACCATTAAC GGAATTC

57

## (2) INFORMATION FOR SEQ ID NO:27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Thr Ser Gly His His His His His His  
 1 5

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 98/06040

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/49 C12N15/62 C07K14/16 A61K39/21

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                                                    | Relevant to claim No. |
|------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X          | WO 94 04686 A (BARSOUM JAMES G.; BIOGEN INC (US); FAWELL STEPHEN E (US); PEPINSKY)<br>3 March 1994<br>see page 54 - page 73<br>---                                                                                                                                                                                    | 1,4,<br>13-15         |
| X          | BODÉUS M ET AL.: "In vitro binding and phosphorylation of human immunodeficiency virus type 1 Nef protein by serine/threonine protein kinase"<br>JOURNAL OF GENERAL VIROLOGY,<br>vol. 76, no. 6, June 1995, pages 1337-1344, XP002092508<br>READING GB<br>see page 1338, left-hand column, paragraph 3<br>---<br>-/-- | 1,5,<br>13-15         |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

5 February 1999

Date of mailing of the international search report

18/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Cupido, M



# INTERNATIONAL SEARCH REPORT

Original Application No

PCT/EP 98/06040

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                    | Relevant to claim No. |
|------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X          | SALFELD J ET AL: "A tripartite HIV-1<br>tat-env-rev fusion protein"<br>EMBO JOURNAL,<br>vol. 9, no. 3, 1 March 1990, pages<br>965-970, XP000113784<br>see the whole document                                                                                                          | 1,4                   |
| X          | AHMED A AZAD ET AL: "Large-scale<br>production and characterization of<br>recombinant human immunodeficiency virus<br>type 1 Nef"<br>JOURNAL OF GENERAL VIROLOGY,<br>vol. 75, no. 3, 1 March 1994, pages<br>651-655, XP000565729<br>see the whole document                            | 1,5,<br>13-15         |
| A          | JANSON H ET AL.: "Protein D, the<br>immunoglobulin D-binding protein of<br>Haemophilus influenzae, is a lipoprotein"<br>INFECTION AND IMMUNITY,<br>vol. 60, no. 4, April 1992, pages<br>1336-1342, XP002092509<br>WASHINGTON US<br>cited in the application<br>see the whole document | 6-8                   |

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/06040

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|-------------------------------------------|---------------------|----------------------------|---------------------|
| WO 9404686 A                              | 03-03-1994          | AT 173016 T                | 15-11-1998          |
|                                           |                     | AU 667244 B                | 14-03-1996          |
|                                           |                     | AU 5083293 A               | 15-03-1994          |
|                                           |                     | CA 2135642 A               | 03-03-1994          |
|                                           |                     | DE 69321962 D              | 10-12-1998          |
|                                           |                     | DE 656950 T                | 14-03-1996          |
|                                           |                     | EP 0656950 A               | 14-06-1995          |
|                                           |                     | ES 2123062 T               | 01-01-1999          |
|                                           |                     | FI 945248 A                | 05-01-1995          |
|                                           |                     | JP 10033186 A              | 10-02-1998          |
|                                           |                     | JP 2702285 B               | 21-01-1998          |
|                                           |                     | JP 7503617 T               | 20-04-1995          |
|                                           |                     | NO 944273 A                | 17-02-1995          |
|                                           |                     | NZ 255831 A                | 24-04-1997          |
|                                           |                     | US 5674980 A               | 07-10-1997          |
|                                           |                     | US 5670617 A               | 23-09-1997          |
|                                           |                     | US 5652122 A               | 29-07-1997          |
|                                           |                     | US 5747641 A               | 05-05-1998          |